

# Maturation of Nineteen Species of Finfish off the Northeast Coast of the United States, 1985-1990

Loretta O'Brien  
Jay Burnett  
Ralph K. Mayo

LIBRARY

JUL 17 1993

National Marine Fisheries Service  
2570 Dole Street  
Honolulu, Hawaii

U.S. Department of Commerce

Library  
NOAA, National Marine Fisheries Service  
SWFC Honolulu Laboratory F/SWC2  
2570 Dole Street  
Honolulu, HI 96822-2396

NOAA  
Tech.  
Rept.  
NMFS  
113  
c.1

## NOAA Technical Reports NMFS

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for their optimum use. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, with the development and enforcement of domestic fisheries regulations, with the surveillance of foreign fishing off U.S. coastal waters, and with the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing services and economic analysis programs and through mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The NOAA Technical Report NMFS series was established in 1983 to replace two subcategories of the Technical Report series: "Special Scientific Report—Fisheries" and "Circular." The series contains the following types of

reports: scientific investigations that document long-term continuing programs of NMFS; intensive scientific reports on studies of restricted scope; papers on applied fishery problems; technical reports of general interest intended to aid conservation and management; reports that review, in considerable detail and at a high technical level, certain broad areas of research; and technical papers originating in economics studies and in management investigations. Since this is a formal series, all submitted papers receive peer review and all papers, once accepted, receive professional editing before publication.

Copies of NOAA Technical Reports NMFS are available free in limited numbers to government agencies, both federal and state. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained from the U.S. Department of Commerce, National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161.

### Recently Published NOAA Technical Reports NMFS

- |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <p><b>100. Marine flora and fauna of the eastern United States: Dicyemida</b>, by Robert B. Short. February 1991, 16 p.</p> <p><b>101. Larvae of nearshore fishes in oceanic waters near Oahu, Hawaii</b>, by Thomas A. Clarke. March 1991, 19 p.</p> <p><b>102. Marine ranching: proceedings of the seventeenth U.S.-Japan meeting on aquaculture; Ise, Mie Prefecture, Japan, 16-18 October 1988</b>, edited by Ralph S. Svrjcek. May 1991, 180 p.</p> <p><b>103. Benthic macrofauna of the New York Bight, 1979-89</b>, by Robert N. Reid, David J. Radosh, Ann B. Frame, and Steven A. Fromm. December 1991, 50 p.</p> <p><b>104. Incidental catch of marine mammals by foreign and joint venture trawl vessels in the U.S. EEZ of the North Pacific, 1973-88</b>, by Michael A. Perez and Thomas R. Loughlin. December 1991, 57 p.</p> <p><b>105. Biology, oceanography, and fisheries of the North Pacific transition zone and subarctic frontal zone</b>, edited by Jerry A. Wetherall. December 1991, 92 p.</p> <p><b>106. Marine ranching: proceedings of the eighteenth U.S.-Japan meeting on aquaculture; Port Ludlow, Washington, 18-19 September 1989</b>, edited by Ralph</p> | <p>S. Svrjcek. February 1992, 136 p.</p> <p><b>107. Field guide to the scarabins (<i>Prionotus</i> and <i>Bel-lator</i>) in the western North Atlantic</b>, by Mike Russell, Mark Grace, and Elmer J. Gutherz. March 1992, 26 p.</p> <p><b>108. Marine debris survey manual</b>, by Christine A. Ribic, Trevor R. Dixon, and Ivan Vining. April 1992, 92 p.</p> <p><b>109. Seasonal climatologies and variability of eastern tropical Pacific surface waters</b>, by Paul C. Fiedler. April 1992, 65 p.</p> <p><b>110. The distribution of Kemp's ridley sea turtles (<i>Lepidochelys kempi</i>) along the Texas coast: an atlas</b>, by Sharon A. Manzella and Jo A. Williams. May 1992, 52 p.</p> <p><b>111. Control of disease in aquaculture: proceedings of the nineteenth U.S.-Japan meeting on aquaculture; Ise, Mie Prefecture, Japan, 29-30 October 1990</b>, edited by Ralph S. Svrjcek. October 1992, 143 p.</p> <p><b>112. Variability of temperature and salinity in the Middle Atlantic Bight and Gulf of Maine</b>, by Robert L. Benway, Jack W. Jossi, Kevin P. Thomas, and Julien R. Goulet. April 1993, 108 p.</p> |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|



NOAA Technical Report NMFS 113

**Maturation of Nineteen Species of Finfish  
off the Northeast Coast of the  
United States, 1985–1990**

Loretta O'Brien  
Jay Burnett  
Ralph K. Mayo

June 1993



U.S. DEPARTMENT OF COMMERCE

Ronald H. Brown, Secretary

National Oceanic and Atmospheric Administration

D. James Baker, Under Secretary for Oceans and Atmosphere

National Marine Fisheries Service

### **Copyright Law**

*Although the contents of these reports have not been copyrighted and may be reprinted entirely, reference to source is appreciated.*

The National Marine Fisheries Service (NMFS) does not approve, recommend, or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

# Contents

|                                                      |           |
|------------------------------------------------------|-----------|
| <b>Abstract</b>                                      | <b>1</b>  |
| <b>Introduction</b>                                  | <b>1</b>  |
| <b>Methods</b>                                       | <b>2</b>  |
| <b>Results</b>                                       | <b>4</b>  |
| Atlantic Cod: <i>Gadus morhua</i>                    | 10        |
| Haddock: <i>Melanogrammus aeglefinus</i>             | 14        |
| Pollock: <i>Pollachius virens</i>                    | 18        |
| White Hake: <i>Urophycis tenuis</i>                  | 20        |
| Red Hake: <i>Urophycis chuss</i>                     | 22        |
| Silver Hake: <i>Merluccius bilinearis</i>            | 26        |
| Redfish: <i>Sebastes fasciatus</i>                   | 30        |
| Scup: <i>Stenotomus chrysops</i>                     | 32        |
| Black Sea Bass: <i>Centropristis striata</i>         | 33        |
| Ocean Pout: <i>Macrozoarces americanus</i>           | 34        |
| Yellowtail Flounder: <i>Pleuronectes ferrugineus</i> | 36        |
| Winter Flounder: <i>Pleuronectes americanus</i>      | 42        |
| American Plaice: <i>Hippoglossoides platessoides</i> | 48        |
| Witch Flounder: <i>Glyptocephalus cynoglossus</i>    | 50        |
| Summer Flounder: <i>Paralichthys dentatus</i>        | 52        |
| Windowpane: <i>Scophthalmus aquosus</i>              | 54        |
| Atlantic Herring: <i>Clupea harengus</i>             | 56        |
| Atlantic Mackerel: <i>Scomber scombrus</i>           | 58        |
| Butterfish: <i>Peprilus triacanthus</i>              | 60        |
| <b>Discussion</b>                                    | <b>62</b> |
| <b>Acknowledgments</b>                               | <b>63</b> |
| <b>Citations</b>                                     | <b>63</b> |



Digitized by the Internet Archive  
in 2025

# Maturation of Nineteen Species of Finfish off the Northeast Coast of the United States, 1985 – 1990

LORETTA O'BRIEN, JAY BURNETT, and RALPH K. MAYO

Woods Hole Laboratory  
Northeast Fisheries Science Center  
Woods Hole, MA 02543

## ABSTRACT

Observations on maturation stages of nineteen species of economically important finfish off the Northeast coast of the USA were analyzed to examine relationships between fish size or age, and maturity. Maturation schedules and median lengths ( $L_{50}$ ) and ages ( $A_{50}$ ) at maturation were derived by fitting the logistic model to the observed proportions. Analyses were generally restricted to observations from 1985 to 1990 obtained during stratified random bottom trawl surveys conducted in spring and autumn by the Northeast Fisheries Science Center and the Commonwealth of Massachusetts Division of Marine Fisheries in waters of the continental shelf from Nova Scotia to Cape Hatteras, North Carolina. Butterfish, *Peprius triacanthus*, attained sexual maturity at the smallest median length (11.4 cm, males) and pollock, *Pollachius virens*, at the highest (41.8 cm, males). Median length at maturity for gadiforms ranged from 22.2 to 41.8 cm. Within the pleuronectiforms, median length at maturity ranged from 19.1 to 30.4 cm. Median lengths for the pelagic and miscellaneous demersal species were in the same ranges as the pleuronectiforms. Butterfish also attained sexual maturity at the youngest median age (0.9 yr, both sexes) whereas redfish, *Sebastes fasciatus*, were the latest to mature (5.5 yr, both sexes). For gadids, the median age at maturity ranged from 1.3 to 2.3 yr. Within the pleuronectiforms, median age at maturity ranged from 1.3 to 4.4 yr and, for pelagic species, from 0.9 to 3.0 yr. Median lengths and ages for many species are lower than those reported in earlier studies of the same general region of the Northwest Atlantic.

## Introduction

Maturation rates, along with rates of growth and mortality, are vital to the characterization of the population dynamics of fish stocks. The age at first reproduction has a strong influence upon the population growth rate (Cole 1954); Beverton and Holt (1959) Beverton (1963, 1987) and Garrod and Horwood (1984) have all noted the interdependence of lifespan, growth, and age at first maturity implicit in the compensatory response of exploited fish stocks to fishing pressure.

The interactions between growth and maturation are varied and complex. Alm (1959) and Roff (1981, 1982) identified size and age as codeterminants in the onset of maturity in fish, but for most species, size was the more important factor. Several studies (Molander 1925; Alm 1959; Jørgensen 1990; O'Brien 1990) have attributed observed reductions in the age at maturity to

increases in growth rates, particularly during the juvenile stage (Godo and Moksness 1987). However, there is some evidence that early maturation is associated with poor growth, i.e. at some point the age effect upon the maturation process overrides the size effect (Elkin 1955; Schaffer and Elson 1975; Mayo et al. 1990). It is also possible that observed changes in median age at maturity values are simply artifacts of changes in growth. For example, a species for which maturation is size-dependent would appear to have lowered its median age at maturity if increased growth caused it to attain the critical size at a younger age. Interpretations of maturity analyses, therefore, should be made carefully and with some consideration of possible changes in growth rates.

Maturity estimates by size or age provide information about trends in maturation as stock sizes and environmental conditions fluctuate, and may be used directly in the evaluation of fishery management programs. Estimates of

median length at maturity are used to evaluate mean retention lengths associated with various commercial gear types. Maturity-at-age data are also employed in the calculation of spawning stock biomass per recruit and maximum spawning potential (Gabriel et al. 1989). Estimates of spawning stock biomass and recruitment provide the framework for calculating the biological reference point,  $F_{rep}$  (Sissenwine and Shepherd 1987), for 13 of the species analyzed in this study which are included in the Northeast Multispecies Fishery Management Plan (New England Fishery Management Council 1985).

In this report, we provide length- and age-specific maturation ogives and estimates of median length and age at maturity for 19 species which have supported important commercial and recreational fisheries off the northeast coast of the United States. Exploitation rates for most of the species analyzed are at historically high levels and stock sizes are near or at historically low levels. Since growth and maturation rates are likely to change as stock sizes decline in abundance (Templeman and Bishop 1979; Sinclair et al. 1980; Bowering 1989), it is probable that growth rates for many of the species included in our study have increased in recent years. In view of these trends, the primary goal of this study is to provide a comprehensive summary of recent maturity information for these species.

## Methods

The data analyzed in this study were collected during stratified random bottom trawl surveys conducted in

spring and autumn by the Northeast Fisheries Science Center (NEFSC), National Marine Fisheries Service (Doubleday 1981; NEFC 1988) and the Massachusetts Division of Marine Fisheries (MDMF) (Azarovitz et al. 1989). Surveys conducted by the NEFSC cover the inshore and offshore areas of the continental shelf from Nova Scotia to Cape Hatteras, North Carolina (Fig. 1). The Massachusetts Division of Marine Fisheries conducts surveys in the inshore waters along the coast of Massachusetts (Fig. 1). Data from MDMF were used only for the inshore stocks of winter flounder, *Pleuronectes americanus*, owing to observed mixing of stocks for this species in the offshore NEFSC survey.

During each survey, finfish were sampled for determination of length (to nearest cm), sex, and maturity stage. Fish with forked tails were measured using fork length (FL); otherwise, all fish were measured using total length (TL). Atlantic herring, *Clupea harengus*, were frozen at sea and measured in the laboratory using total length. Black sea bass, *Centropomus striata*, were also measured using total length, excluding the caudal filament. The criteria for seven stages of maturity classification are presented in Table 1 as similarly described by Burnett et al. (1989). Fish were also sampled for age determination by removing either scales or otoliths and were analyzed in the laboratory by using species-specific ageing methods as described by Penttila and Dery (1988).

Nineteen species, comprising twenty-nine stocks, were analyzed to determine the median length ( $L_{50}$ ) at maturity; fourteen species, comprising twenty-two stocks, were analyzed to determine median age ( $A_{50}$ ) at maturity.




Figure 1

Areas of the continental shelf from Nova Scotia to Cape Hatteras, North Carolina surveyed by the Northeast Fisheries Science Center and the Massachusetts Division of Marine Fisheries spring and autumn bottom trawl surveys.

ity. Maturity classifications were considered as either immature or mature by combining all stages other than immature. Length and age based analyses were generally restricted to the period 1985–1989 or 1990 depending upon the availability of data (Table 2). Exceptions were redfish, *Sebastes fasciatus*, and red hake, *Urophycis chuss*, for which recent age data were not available. Stock areas (Fig. 1) were delineated by NEFSC bottom trawl survey strata (Table 2) as defined by current stock

structure (NEFSC 1991). Either spring or autumn data collected closest to the time of spawning were used in the analyses for each species (Table 2). This was done in an attempt to minimize errors in maturity classifications, especially in differentiating between immature and resting gonads (Halliday 1987).

The observed proportion of fish mature at length and age was calculated for each species by stock area, year, season, and sex. Immature fish not identified by

Table 1

Current maturity staging criteria used during Northeast Fisheries Science Center bottom trawl surveys.

| Stage            | Code | Description and criteria                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
|------------------|------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Females</b>   |      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Immature         | I    | Ovary paired, tube-like, small relative to body cavity; colorless to pink jelly-like tissue, no visible eggs; thin, transparent outer membrane.<br>Butterfish, <i>Peprilus triacanthus</i> : ovary paired and flattened, ovary wall striated or wrinkled.<br>Ocean pout, <i>Macrozoarces americanus</i> : ovary <b>not</b> paired.                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Developing       | D    | Ovaries large, occupying up to 2/3 of body cavity; blood vessels prominent when present; ovary appears granular as yellow to orange yolked eggs develop. A mixture of less than 50% yolked and hydrated eggs denotes <b>D</b> .                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Ripe             | R    | Ovaries large, may fill entire body cavity; a mixture of 50% or more hydrated eggs denotes <b>R</b> ; transparent ovary wall.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Eyed             | E    | Redfish <i>Sebastes fasciatus</i> only: ovary large, robust, transparent membrane with dark spotted eggs containing developing eyed-stage larvae.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Ripe and Running | U    | Eggs flow from vent with little or no pressure to abdomen.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Spent            | S    | Ovaries flaccid, sac-like, similar in size to ripe ovaries; color red to purple; ovary wall thickened, cloudy and translucent; some hydrated or yolked eggs may adhere to ovary wall.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Resting          | T    | Ovaries smaller than ripe ovaries, but larger than immature. Interior jelly-like, no visible eggs.<br>Haddock, <i>Melanogrammus aeglefinus</i> : ovaries small relative to body cavity, wall thick and tough with purplish membrane, blood vessels less prominent and purplish.<br>Pleuronectiformes: ovaries larger relative to the body cavity than in gadids, interior yellow or orange. Ovary wall cloudy or translucent and thicker and tougher than in ripe ovaries.                                                                                                                                                                                                                                                                                                                            |
| <b>Males</b>     |      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Immature         | I    | Testes small relative to body cavity, colorless to gray and translucent.<br>Butterfish, <i>Peprilus triacanthus</i> : similar to ovary, but with a smooth wall.<br>Pleuronectiformes: testis located posterior of body cavity; triangular shape with sharp anterior edge.<br>Winter flounder, <i>Pleuronectes americanus</i> : posterior portion of testis extends further back similar to ovary but the shape is different: the endpoint of the testis is rounded, and more flattened as it extends posteriorly.<br>Gadids: testes narrow, lobed and elongated, resembles crimped ribbon.<br>Atlantic herring, <i>Clupea harengus</i> , mackerel <i>Scomber scombrus</i> , scup, <i>Stenotomus chrysops</i> : similar shape as ovary, but there is an edge to the testis vs. being rounded in ovary. |
| Developing       | D    | Testes large, grey to off-white, firm consistency with very little or no milt present.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Ripe             | R    | Testes larger than <b>D</b> , chalk white, consistency mostly liquid. Milt flows easily when testis dissected.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Ripe and Running | U    | Chalk white milt flows easily from the vent with little or no pressure on abdomen. Once dissected, milt flows easily.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Spent            | S    | Testes flaccid, may contain residual milt, less robust than <b>R</b> . Edges or other parts of the testes starting to turn grey as milt recedes.<br>Gadiformes: edges of lobes reddish to brown or grey as milt recedes.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Resting          | T    | Testes shrunken in size relative to <b>R</b> . Color is yellow, brown, or grey with little or no milt.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |

sex were assumed to have a 1:1 sex ratio and equal catchability and were included in the analyses by adding the number of unsexed fish at each length to both female and male data sets. As a result, the sample size increased by less than 1% (*U. tenuis*) to 24% (*Stenotomus chrysops*) in 27 of the 29 stock analyses. No other adjustments were made to the observations.

The proportion of fish mature at length and age was estimated by fitting the logistic model to the observed proportions mature at length and age. The form of the logistic was

$$P = \frac{1}{1 + e^{-(\alpha + \beta x)}}, \quad (1)$$

where  $P$  = proportion mature at age or length  
 $x$  = either length or age, and  
 $\alpha, \beta$  = model parameters to be estimated.

Logistic regressions (Dixon 1981) were performed on each species stock, by sex, to obtain the model parameters. Median length or age at maturity was calculated from the regression coefficients as  $[-a/b]$ .

The generally low sample sizes for each sex within a stock area did not allow for calculation of reliable annual estimates. Thus, years with similar values of  $L_{50}$  and  $A_{50}$  that also had relatively stable stock sizes were pooled to achieve the objective of obtaining a representative maturity schedule reflecting current stock conditions. As a result, one or two years of data were excluded for five species and nine analyses. The combined data were then analyzed using the same methods as described above.

The approximate variance for large samples (Ashton 1972) was calculated for given proportions mature ( $y$ ) at length or age as

$$s^2(y) = \frac{1}{b^2} \left[ s^2(a) + \frac{a^2}{b^2} s^2(b) - \frac{2a}{b} \text{cov}(a, b) \right]. \quad (2)$$

The computational equation for this formula is

$$s^2(y) = \frac{1}{b^2} \left[ s^2(\hat{a}) + (y - \bar{x})^2 s^2(b) \right], \quad (3)$$

where  $a, b$  = estimates of  $\alpha$  and  $\beta$

$y$  = length or age at maturity for a given proportion

$s^2(\hat{a}) = 1 / \sum (nPQ)$

$P$  = probability of a response

$Q = 1 - P$

$n$  = number of fish at length or age  $x$

$\bar{x} = \sum nPQx / \sum nPQ$

$s(b)$  = standard error of  $b$ .

Since exact confidence limits for the statistic  $y$  are not available, approximate 95% confidence limits were calculated (Zar 1984) at increments of 0.05 for proportions mature at length and at whole ages for proportions mature at age. The  $t$ -distribution was used at an alpha level of 0.05 for varying degrees of freedom (df) with the standard error of  $y$  as

$$y \pm t_{0.05, df} s(y). \quad (4)$$

## Results

Butterfish, *Peprilus triacanthus*, attained sexual maturity at the smallest median length (11.4 cm, males) and pollock, *Pollachius virens*, at the highest (41.8 cm, males) (Table 3). Median length at maturity for the gadids ranged from 22.2 cm (red hake, *U. chuss*, males) to 41.8 cm (pollock, *P. virens*, males). Within the pleuronectiforms, median length at maturity ranged from 19.1 cm (yellowtail flounder, *Pleuronectes ferrugineus*, males) to 30.4 cm (witch flounder, *Glyptocephalus cynoglossus*, females). Median lengths for the pelagic and miscellaneous demersal species were in the same ranges as the pleuronectiforms.

Butterfish, *Peprilus triacanthus*, also attained sexual maturity at the youngest median age (0.9 yr, both sexes) while redfish, *Sebastes fasciatus*, were the latest to mature (5.5 yr, both sexes) (Table 4). For gadids, the median age at maturity ranged from 1.3 yr (haddock, *Melanogrammus aeglefinus*, males) to 2.3 yr (cod, *Gadus morhua*, and pollock, *Pollachius virens*, males). Within the pleuronectiforms, median age at maturity ranged from 1.3 yr (yellowtail flounder, *Pleuronectes ferrugineus*, males) to 4.4 yr (witch flounder, *G. cynoglossus*, females) and, for pelagic species, from 0.9 yr (butterfish, *Peprilus triacanthus*) to 3.0 yr (Atlantic herring, *Clupea harengus*, females). Median lengths and ages for many species are lower than those reported in earlier studies of the same general region of the Northwest Atlantic.

Median lengths vs. median ages at maturation for the gadiforms and pleuronectiforms are plotted in Figure 2. For gadiforms,  $L_{50}$  and  $A_{50}$  estimates extend over a wide range of lengths within a relatively narrow age span. Pleuronectiforms, however, exhibit an opposite pattern, with estimates extending over a very wide range of ages within a narrow length span. This suggests that, during the study period, the maturation process was age-dependent for the gadiforms and size-dependent for the pleuronectiforms.

In each of the following species sections, tables of the proportion of fish mature at length are presented in 0.05 increments and tables of proportions of fish mature at age are presented in whole ages.

Table 2

Species by stock area, spawning season (Smith 1985<sup>a</sup>), survey season, Northeast Fisheries Science Center offshore strata sets, and years included in each species stock analysis (GB=Georges Bank, NGB=Northern Georges Bank, SGB=Southern Georges Bank, GM=Gulf of Maine, CC=Cape Cod, SNE=Southern New England, MA=Middle Atlantic, N=North, S=South).

| Species                             | Area    | Spawning season<br>(peak) | Survey<br>Analyzed  | Strata set                                                            | Years     |
|-------------------------------------|---------|---------------------------|---------------------|-----------------------------------------------------------------------|-----------|
| Atlantic Cod                        | GB      | Nov-May (Feb-Mar)         | Spring              | 13-25                                                                 | 1985-1990 |
| <i>Gadus morhua</i>                 | GM      | Nov-May (Mar-May)         | Spring              | 26-30, 36-40                                                          | 1985-1990 |
| Haddock                             | GB      | Jan-May (Mar-Apr)         | Spring              | 13-25, 29-30                                                          | 1985-1989 |
| <i>Melanogrammus aeglefinus</i>     | GM      | Feb-May (Mar-Apr)         | Spring              | 26-28, 36-40                                                          | 1985-1989 |
| Pollock                             | GB-GM   | Oct-Mar (Dec-Jan)         | Autumn              | 13-34, 351,                                                           | 1986-1988 |
| <i>Pollachius virens</i>            |         |                           |                     | 36-40                                                                 |           |
| White hake                          | GB-GM   | Oct-May                   | Spring              | 21-30, 33, 34                                                         | 1987-1989 |
| <i>Urophycis tenuis</i>             |         |                           |                     | 351, 36-40                                                            |           |
| Red hake                            | GM-NGB  | May-Sep (Jun-Jul)         | Spring              | 20-30, 36-40                                                          | 1982-1985 |
| <i>Urophycis chuss</i>              | SGB-MA  | Mar-Oct (Aug-Sep)         | Spring              | 1-19, 61-76;<br>1-46 <sup>b</sup> , 52 <sup>b</sup> , 55 <sup>b</sup> |           |
| Silver hake                         | GM-NGB  | Apr-Oct (May-Aug)         | Spring              | 20-30, 36-40                                                          | 1985-1989 |
| <i>Merluccius bilinearis</i>        | SGB-MA  | Apr-Oct (May-Sep)         | Spring              | 1-19, 61-76;<br>1-46 <sup>b</sup> , 52 <sup>b</sup> , 55 <sup>b</sup> |           |
| Redfish                             | GM      | Apr-Aug (Jun-Jul)         | Spring <sup>c</sup> | 24, 26-30                                                             | 1975-1980 |
| <i>Sebastes fasciatus</i>           |         |                           | Autumn <sup>d</sup> | 36-40                                                                 |           |
| Scup                                | All     | May-Aug (May-Jun)         | Spring              | 2-3, 61-63,                                                           | 1985      |
| <i>Stenotomus chrysops</i>          |         |                           |                     | 65-67, 70-71, 74-75                                                   | 1987-1990 |
| Black sea bass                      | All     | Apr-Oct (Jun)             | Spring              | 1-12, 25, 61-76;                                                      | 1985-1990 |
| <i>Centropristis striata</i>        |         |                           |                     | 1-61 <sup>b</sup>                                                     |           |
| Ocean pout                          | SNE     | Sep-Oct (Oct)             | Spring              | 1-12, 61-76;                                                          | 1985-1990 |
| <i>Macrozoarces americanus</i>      |         |                           |                     | 1-52 <sup>b</sup>                                                     |           |
| Yellowtail flounder                 | GM      | Sep-Oct (Sep)             | Spring              | 26-40; 57-66 <sup>b</sup>                                             | 1985-1990 |
| <i>Pleuronectes ferrugineus</i>     | GB      | Apr-Aug (May-Jun)         | Spring              | 13-21                                                                 | 1985-1990 |
|                                     | CC      | Apr-Aug (May-Jun)         | Spring              | 24-26                                                                 |           |
|                                     | SNE     | Apr-Aug (May-Jun)         | Spring              | 5, 6, 9, 10                                                           |           |
| Winter flounder                     | GB      | Mar-May (Apr)             | Spring              | 13-22                                                                 | 1985-1989 |
| <i>Pleuronectes americanus</i>      | N of CC | Apr-May (Apr)             | Spring              | 25-36 <sup>e</sup>                                                    |           |
|                                     | S of CC | Feb-May (Feb-Mar)         | Spring              | 11-17 <sup>e</sup>                                                    |           |
| American plaice                     | GM      | Feb-Jun (Apr-May)         | Spring              | 26-30, 36-40                                                          | 1986-1990 |
| <i>Hippoglossoides platessoides</i> |         |                           |                     |                                                                       |           |
| Witch flounder                      | GM      | Apr-Aug (May-Jun)         | Spring              | 22-30, 33-34,                                                         | 1986-1990 |
| <i>Glyptocephalus cynoglossus</i>   |         |                           |                     | 351, 36-40                                                            |           |
| Summer flounder                     | All     | Aug-Dec (Oct-Nov)         | Autumn              | 1-12, 45-46,                                                          | 1985-1989 |
| <i>Paralichthys dentatus</i>        |         |                           |                     | 52, 55-76; 1-41 <sup>b</sup>                                          |           |
| Windowpane                          | GB      | Jun-Oct (Jul-Aug)         | Spring              | 13-21                                                                 | 1985-1990 |
| <i>Scophthalmus aquosus</i>         | SNE     | May-Nov (Sep)             | Spring              | 5-12                                                                  |           |
| Atlantic herring                    | SNE,GM  | Aug-Dec (Sep-Oct)         | Autumn              | 1-12, 24-30,                                                          | 1987-1989 |
| <i>Clupea harengus</i>              |         |                           |                     | 33-34, 351,<br>36-40, 61-76                                           |           |
| Atlantic mackerel                   | All     | Apr-Jul (Apr-Jun)         | Spring              | 1-34, 351,                                                            | 1987-1989 |
| <i>Scomber scombrus</i>             |         |                           |                     | 36-76; 1-81 <sup>b</sup>                                              |           |
| Butterfish                          | All     | May-Sep (Jun-Jul)         | Spring              | 1-25, 61-76;                                                          | 1986-1989 |
| <i>Peprilus triacanthus</i>         |         |                           |                     | 1-46 <sup>b</sup>                                                     |           |

<sup>a</sup> W.G. Smith. 1985. Temporal and spatial spawning patterns of the principal species of fish and invertebrates in the Georges Bank region. NMFS/NEFC, Sandy Hook Laboratory, Lab. Ref. Doc. No. 85-4, 35 p.

<sup>b</sup> Northeast Fisheries Science Center inshore strata.

<sup>c</sup> females only.

<sup>d</sup> males only.

<sup>e</sup> Massachusetts Division of Marine Fisheries strata.

Table 3

Median length (cm) at maturity ( $L_{50}$ ), parameter values  $a$  and  $b$  from the logistic regression, standard error (SE) of  $a$  and  $b$ , and the number of females (F) and males (M) of nineteen species of finfish sampled off the northeast coast of the United States.

| Species                                      | Sex | $a$     | SE $a$ | $b$   | SE $b$ | $L_{50}$ | $n$  |
|----------------------------------------------|-----|---------|--------|-------|--------|----------|------|
| Atlantic cod, <i>Gadus morhua</i>            |     |         |        |       |        |          |      |
| 1985–1990                                    |     |         |        |       |        |          |      |
| Georges Bank                                 | F   | -4.932  | 0.429  | 0.127 | 0.009  | 38.8     | 983  |
|                                              | M   | -5.688  | 0.464  | 0.139 | 0.010  | 41.0     | 1034 |
| Gulf of Maine                                | F   | -5.500  | 0.653  | 0.171 | 0.018  | 32.1     | 411  |
|                                              | M   | -5.048  | 0.554  | 0.140 | 0.014  | 36.0     | 394  |
| Haddock, <i>Melanogrammus aeglefinus</i>     |     |         |        |       |        |          |      |
| 1985–1989                                    |     |         |        |       |        |          |      |
| Georges Bank                                 | F   | -6.252  | 0.765  | 0.210 | 0.022  | 29.7     | 580  |
|                                              | M   | -5.064  | 0.738  | 0.189 | 0.022  | 26.8     | 478  |
| Gulf of Maine                                | F   | -18.605 | 11.78  | 0.539 | 0.314  | 34.5     | 69   |
|                                              | M   | -13.111 | 5.432  | 0.374 | 0.146  | 35.0     | 49   |
| Pollock, <i>Pollachius virens</i>            |     |         |        |       |        |          |      |
| 1986–1988                                    |     |         |        |       |        |          |      |
|                                              | F   | -6.004  | 1.120  | 0.154 | 0.027  | 39.1     | 154  |
|                                              | M   | -6.788  | 1.158  | 0.162 | 0.028  | 41.8     | 143  |
| White hake, <i>Urophycis tenuis</i>          |     |         |        |       |        |          |      |
| 1987–1989                                    |     |         |        |       |        |          |      |
|                                              | F   | -4.808  | 0.538  | 0.137 | 0.014  | 35.1     | 455  |
|                                              | M   | -7.790  | 0.894  | 0.238 | 0.026  | 32.7     | 346  |
| Red hake, <i>Urophycis chuss</i>             |     |         |        |       |        |          |      |
| 1982–1985                                    |     |         |        |       |        |          |      |
| Gulf of Maine - Northern                     | F   | -13.560 | 1.665  | 0.504 | 0.058  | 26.9     | 667  |
|                                              | M   | -12.073 | 1.846  | 0.543 | 0.075  | 22.2     | 595  |
| Southern Georges Bank-Middle                 | F   | -12.767 | 1.165  | 0.509 | 0.043  | 25.1     | 1020 |
|                                              | M   | -11.614 | 1.138  | 0.488 | 0.043  | 23.8     | 753  |
| Atlantic                                     |     |         |        |       |        |          |      |
| Silver hake, <i>Merluccius bilinearis</i>    |     |         |        |       |        |          |      |
| 1985–1989                                    |     |         |        |       |        |          |      |
| Gulf of Maine - Northern                     | F   | -11.963 | 0.687  | 0.519 | 0.029  | 23.1     | 1410 |
|                                              | M   | -14.369 | 0.938  | 0.646 | 0.041  | 22.3     | 1026 |
| Southern Georges Bank-Middle                 | F   | -10.753 | 0.537  | 0.464 | 0.021  | 23.2     | 2429 |
|                                              | M   | -14.263 | 0.890  | 0.629 | 0.037  | 22.7     | 1612 |
| Atlantic                                     |     |         |        |       |        |          |      |
| Redfish, <i>Sebastes fasciatus</i>           |     |         |        |       |        |          |      |
| 1975–1980                                    |     |         |        |       |        |          |      |
|                                              | F   | -13.240 | 1.017  | 0.595 | 0.044  | 22.3     | 1457 |
|                                              | M   | -6.929  | 0.476  | 0.331 | 0.020  | 20.9     | 1526 |
| Scup, <i>Stenotomus chrysops</i>             |     |         |        |       |        |          |      |
| 1985, 1987–1990                              |     |         |        |       |        |          |      |
|                                              | F   | -14.071 | 1.944  | 0.907 | 0.122  | 15.5     | 284  |
|                                              | M   | -12.491 | 1.797  | 0.801 | 0.112  | 15.6     | 232  |
| Black sea bass, <i>Centropristis striata</i> |     |         |        |       |        |          |      |
| 1985–1990                                    |     |         |        |       |        |          |      |
|                                              | F   | -4.691  | 0.535  | 0.245 | 0.024  | 19.1     | 561  |
|                                              | M   | -6.841  | 0.957  | 0.360 | 0.045  | 19.0     | 348  |
| Ocean Pout, <i>Macrozoarces americanus</i>   |     |         |        |       |        |          |      |
| 1985–1990                                    |     |         |        |       |        |          |      |
| Gulf of Maine                                | F   | -5.101  | 2.103  | 0.195 | 0.065  | 26.2     | 88   |
|                                              | M   | -4.455  | 2.343  | 0.147 | 0.057  | 30.3     | 55   |

Table 3 (Continued)

Median length (cm) at maturity ( $L_{50}$ ), parameter values  $a$  and  $b$  from the logistic regression, standard error (SE) of  $a$  and  $b$ , and the number of females (F) and males (M) of nineteen species of finfish sampled off the northeast coast of the United States.

| Species                                                           | Sex | $a$     | SE $a$ | $b$   | SE $b$ | $L_{50}$ | $n$ |
|-------------------------------------------------------------------|-----|---------|--------|-------|--------|----------|-----|
| Southern New England                                              | F   | -6.515  | 1.584  | 0.208 | 0.040  | 31.3     | 197 |
|                                                                   | M   | -5.920  | 1.204  | 0.186 | 0.032  | 31.9     | 205 |
| Yellowtail flounder, <i>Pleuronectes ferrugineus</i><br>1985-1990 |     |         |        |       |        |          |     |
| Georges Bank                                                      | F   | -15.136 | 5.207  | 0.587 | 0.181  | 25.8     | 167 |
|                                                                   | M   | -8.145  | 2.776  | 0.381 | 0.098  | 21.4     | 271 |
| Cape Cod                                                          | F   | -11.738 | 1.951  | 0.430 | 0.071  | 27.3     | 140 |
|                                                                   | M   | -10.103 | 1.577  | 0.377 | 0.058  | 26.8     | 165 |
| Southern New England                                              | F   | -11.366 | 2.833  | 0.445 | 0.098  | 25.5     | 204 |
|                                                                   | M   | -7.331  | 1.734  | 0.375 | 0.066  | 19.6     | 367 |
| Winter flounder, <i>Pleuronectes americanus</i><br>1985-1989      |     |         |        |       |        |          |     |
| Georges Bank                                                      | F   | -10.855 | 2.245  | 0.436 | 0.084  | 24.9     | 204 |
|                                                                   | M   | -7.724  | 1.551  | 0.301 | 0.054  | 25.6     | 174 |
| North of Cape Cod                                                 | F   | -26.646 | 3.661  | 0.896 | 0.121  | 29.7     | 320 |
|                                                                   | M   | -16.906 | 2.416  | 0.613 | 0.087  | 27.6     | 215 |
| South of Cape Cod                                                 | F   | -11.895 | 1.320  | 0.431 | 0.046  | 27.6     | 398 |
|                                                                   | M   | -17.099 | 2.130  | 0.590 | 0.074  | 29.0     | 301 |
| American plaice, <i>Hippoglossoides platessoides</i><br>1986-1990 |     |         |        |       |        |          |     |
|                                                                   | F   | -11.255 | 0.944  | 0.420 | 0.034  | 26.8     | 714 |
|                                                                   | M   | -7.414  | 0.612  | 0.336 | 0.026  | 22.1     | 633 |
| Witch flounder, <i>Glyptocephalus cynoglossus</i><br>1986-1990    |     |         |        |       |        |          |     |
|                                                                   | F   | -11.930 | 2.698  | 0.392 | 0.084  | 30.4     | 189 |
|                                                                   | M   | -9.405  | 2.239  | 0.371 | 0.078  | 25.3     | 233 |
| Summer flounder, <i>Paralichthys dentatus</i><br>1985-1989        |     |         |        |       |        |          |     |
|                                                                   | F   | -8.574  | 1.064  | 0.306 | 0.034  | 28.0     | 378 |
|                                                                   | M   | -7.636  | 0.852  | 0.307 | 0.030  | 24.9     | 497 |
| Windowpane, <i>Scophthalmus aquosus</i><br>1985-1990              |     |         |        |       |        |          |     |
| Georges Bank                                                      | F   | -15.260 | 3.777  | 0.678 | 0.150  | 22.5     | 343 |
|                                                                   | M   | -15.670 | 3.535  | 0.704 | 0.145  | 22.2     | 199 |
| Southern New England                                              | F   | -88.592 | 72.23  | 4.185 | 3.441  | 21.2     | 143 |
|                                                                   | M   | -19.595 | 4.553  | 0.910 | 0.201  | 21.5     | 133 |
| Atlantic herring, <i>Clupea harengus</i><br>1987-1989             |     |         |        |       |        |          |     |
|                                                                   | F   | -28.484 | 3.955  | 1.123 | 0.151  | 25.4     | 359 |
|                                                                   | M   | -40.461 | 6.218  | 1.597 | 0.243  | 25.3     | 307 |
| Atlantic mackerel, <i>Scomber scombrus</i><br>1987-1989           |     |         |        |       |        |          |     |
|                                                                   | F   | -16.475 | 1.601  | 0.641 | 0.060  | 25.7     | 697 |
|                                                                   | M   | -17.102 | 1.516  | 0.658 | 0.056  | 26.0     | 770 |
| Butterfish, <i>Peprilus triacanthus</i><br>1986-1989              |     |         |        |       |        |          |     |
|                                                                   | F   | -14.495 | 1.939  | 1.213 | 0.157  | 12.0     | 333 |
|                                                                   | M   | -11.029 | 1.333  | 0.968 | 0.112  | 11.4     | 341 |

Table 4

Median age (yrs) at maturity ( $A_{50}$ ), parameter values  $a$  and  $b$  from the logistic regression, standard error (SE) of  $a$  and  $b$ , and the number of females (F) and males (M) of fourteen species of finfish sampled off the northeast coast of the United States.

| Species                                              | Sex | $a$     | SE $a$ | $b$   | SE $b$ | $A_{50}$ | $n$  |
|------------------------------------------------------|-----|---------|--------|-------|--------|----------|------|
| Atlantic cod, <i>Gadus morhua</i>                    |     |         |        |       |        |          |      |
| 1985–1990                                            |     |         |        |       |        |          |      |
| Georges Bank                                         | F   | -2.842  | 0.301  | 1.626 | 0.131  | 1.7      | 970  |
|                                                      | M   | -3.258  | 0.306  | 1.725 | 0.129  | 1.9      | 1021 |
| Gulf of Maine                                        | F   | -4.690  | 0.571  | 2.225 | 0.233  | 2.1      | 409  |
|                                                      | M   | -3.828  | 0.451  | 1.651 | 0.173  | 2.3      | 392  |
| Haddock, <i>Melanogrammus aeglefinus</i>             |     |         |        |       |        |          |      |
| 1985–1989                                            |     |         |        |       |        |          |      |
| Georges Bank                                         | F   | -3.283  | 0.482  | 2.233 | 0.250  | 1.5      | 580  |
|                                                      | M   | -2.784  | 0.494  | 2.189 | 0.275  | 1.3      | 478  |
| Gulf of Maine                                        | F   | -4.423  | 1.996  | 2.504 | 0.792  | 1.8      | 69   |
|                                                      | M   | -4.190  | 1.915  | 1.963 | 0.698  | 2.1      | 49   |
| Pollock, <i>Pollachius virens</i>                    |     |         |        |       |        |          |      |
| 1986–1988                                            |     |         |        |       |        |          |      |
|                                                      | F   | -2.346  | 0.554  | 1.173 | 0.221  | 2.0      | 154  |
|                                                      | M   | -4.415  | 0.823  | 1.897 | 0.347  | 2.3      | 143  |
| White hake, <i>Urophycis tenuis</i>                  |     |         |        |       |        |          |      |
| 1987–1989                                            |     |         |        |       |        |          |      |
|                                                      | F   | -2.441  | 0.324  | 1.751 | 0.180  | 1.4      | 455  |
|                                                      | M   | -3.304  | 0.418  | 2.400 | 0.262  | 1.4      | 346  |
| Red hake, <i>Urophycis chuss</i>                     |     |         |        |       |        |          |      |
| 1982–1985                                            |     |         |        |       |        |          |      |
| Gulf of Maine - Northern                             | F   | -7.342  | 1.167  | 4.166 | 0.587  | 1.8      | 667  |
| Georges Bank                                         | M   | -5.725  | 0.846  | 3.994 | 0.473  | 1.4      | 595  |
| Southern Georges                                     |     |         |        |       |        |          |      |
| Bank-Middle                                          | F   | -5.554  | 0.485  | 3.205 | 0.251  | 1.7      | 1020 |
| Atlantic                                             | M   | -4.957  | 0.452  | 2.799 | 0.235  | 1.8      | 753  |
| Silver hake, <i>Merluccius bilinearis</i>            |     |         |        |       |        |          |      |
| 1985–1989                                            |     |         |        |       |        |          |      |
| Gulf of Maine - Northern                             | F   | -5.825  | 0.323  | 3.342 | 0.175  | 1.7      | 1410 |
| Georges Bank                                         | M   | -6.349  | 0.374  | 3.710 | 0.213  | 1.7      | 1026 |
| Southern Georges                                     |     |         |        |       |        |          |      |
| Bank-Middle                                          | F   | -5.748  | 0.272  | 3.599 | 0.152  | 1.6      | 2429 |
| Atlantic                                             | M   | -6.176  | 0.337  | 3.746 | 0.187  | 1.6      | 1612 |
| Redfish, <i>Sebastes fasciatus</i>                   |     |         |        |       |        |          |      |
| 1975–1980                                            |     |         |        |       |        |          |      |
|                                                      | F   | -6.455  | 0.563  | 1.172 | 0.096  | 5.5      | 1457 |
|                                                      | M   | -3.336  | 0.331  | 0.602 | 0.050  | 5.5      | 1526 |
| Yellowtail flounder, <i>Pleuronectes ferrugineus</i> |     |         |        |       |        |          |      |
| 1985–1990                                            |     |         |        |       |        |          |      |
| Georges Bank                                         | F   | -17.150 | 29.78  | 9.578 | 14.89  | 1.8      | 167  |
|                                                      | M   | -5.152  | 2.219  | 3.866 | 1.131  | 1.3      | 271  |
| Cape Cod                                             | F   | -10.355 | 1.572  | 3.933 | 0.595  | 2.6      | 140  |
|                                                      | M   | -7.388  | 1.117  | 2.874 | 0.427  | 2.6      | 165  |
| Southern New England                                 | F   | -4.886  | 1.049  | 2.967 | 0.529  | 1.6      | 204  |
|                                                      | M   | -18.942 | 25.49  | 10.72 | 12.75  | 1.8      | 367  |
| Winter flounder, <i>Pleuronectes americanus</i>      |     |         |        |       |        |          |      |
| 1985–1989                                            |     |         |        |       |        |          |      |
| Georges Bank                                         | F   | -7.164  | 2.075  | 3.822 | 1.011  | 1.9      | 204  |
|                                                      | M   | -4.728  | 1.155  | 2.481 | 0.523  | 1.9      | 174  |
| North of Cape Cod                                    | F   | -11.903 | 1.259  | 3.423 | 0.355  | 3.5      | 320  |
|                                                      | M   | -8.568  | 1.221  | 2.627 | 0.384  | 3.3      | 215  |
| South of Cape Cod                                    | F   | -8.286  | 1.027  | 2.801 | 0.334  | 3.0      | 398  |
|                                                      | M   | -7.706  | 0.927  | 2.317 | 0.285  | 3.3      | 301  |

Table 4 (Continued)

Median age (yrs) at maturity ( $A_{50}$ ), parameter values  $a$  and  $b$  from the logistic regression, standard error (SE) of  $a$  and  $b$ , and the number of females (F) and males (M) of fourteen species of finfish sampled off the northeast coast of the United States.

| Species                                                           | Sex | $a$     | SE $a$ | $b$   | SE $b$ | $A_{50}$ | $n$ |
|-------------------------------------------------------------------|-----|---------|--------|-------|--------|----------|-----|
| American plaice, <i>Hippoglossoides platessoides</i><br>1986–1990 | F   | -7.419  | 0.584  | 2.089 | 0.160  | 3.6      | 714 |
|                                                                   | M   | -4.975  | 0.413  | 1.659 | 0.132  | 3.0      | 633 |
| Witch flounder, <i>Glyptocephalus cynoglossus</i><br>1986–1990    | F   | -6.957  | 1.562  | 1.591 | 0.327  | 4.4      | 189 |
|                                                                   | M   | -8.468  | 2.284  | 2.340 | 0.563  | 3.6      | 233 |
| Atlantic herring, <i>Clupea harengus</i><br>1987–1989             | F   | -14.503 | 3.101  | 4.804 | 1.018  | 3.0      | 359 |
|                                                                   | M   | -11.480 | 2.245  | 3.938 | 0.726  | 2.9      | 307 |
| Atlantic mackerel, <i>Scomber scombrus</i><br>1987–1989           | F   | -8.499  | 1.426  | 4.511 | 0.722  | 1.9      | 697 |
|                                                                   | M   | -6.068  | 0.718  | 3.199 | 0.366  | 1.9      | 770 |
| Butterfish, <i>Peprilus triacanthus</i><br>1986–1989              | F   | -2.608  | 0.590  | 3.001 | 0.521  | 0.9      | 333 |
|                                                                   | M   | -2.553  | 0.574  | 2.909 | 0.521  | 0.9      | 341 |



Figure 2

Median length ( $L_{50}$ ) and median age ( $A_{50}$ ) at maturity for female and male Gadiformes and Pleuronectiformes.

## Atlantic Cod: *Gadus morhua*

Atlantic cod are distributed in the northwest Atlantic from Port Burwell, West Greenland to Cape Lookout, North Carolina (Serchuk and Wood 1979<sup>1</sup>). Within the New England area, four distinct stocks are recognized (Wise 1963): Georges Bank, Gulf of Maine, Southern New England, and the New Jersey coastal cod. The spawning season is from November to May; peak spawning on Georges Bank occurs during February and March and in the Gulf of Maine from March to May (Table 2). Total commercial landings of Atlantic cod from Georges Bank declined from a peak of about 64,000 metric tons (t) in 1982 to a low of about 28,000 t in 1986 and have increased to about 37,000 t in 1989 (NEFSC 1991). Landings from the Gulf of Maine region declined from a peak of about 15,100 t in 1983 to a low of about 8,300

t in 1987 and have recently increased to about 13,000 t in 1989 (NEFSC 1991).

Median length at maturity for female and male Atlantic cod, sampled from Georges Bank during the

<sup>1</sup> F.M. Serchuk and P.W. Wood. 1979. Review and status of the Southern New England - Middle Atlantic Atlantic cod, *Gadus morhua*, populations. NEFC, Woods Hole Laboratory, Lab. Ref. Doc. 79-37, 71 p.




**Figure 3**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of Georges Bank female ( $n=983$ ) and male ( $n=1034$ ) Atlantic cod, *Gadus morhua*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 5**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Georges Bank female ( $n=983$ ) and male ( $n=1034$ ) Atlantic cod, *Gadus morhua*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 15.6           | 11.2                    | 20.1  |
| 0.10                 | 21.5           | 17.8                    | 25.2  |
| 0.15                 | 25.2           | 21.9                    | 28.4  |
| 0.20                 | 27.9           | 25.0                    | 30.8  |
| 0.25                 | 30.2           | 27.6                    | 32.8  |
| 0.30                 | 32.2           | 29.8                    | 34.6  |
| 0.35                 | 34.0           | 31.7                    | 36.2  |
| 0.40                 | 35.6           | 33.6                    | 37.7  |
| 0.45                 | 37.2           | 35.3                    | 39.2  |
| 0.50                 | 38.8           | 37.0                    | 40.6  |
| 0.55                 | 40.4           | 38.7                    | 42.1  |
| 0.60                 | 42.0           | 40.4                    | 43.7  |
| 0.65                 | 43.7           | 42.1                    | 45.3  |
| 0.70                 | 45.5           | 43.9                    | 47.1  |
| 0.75                 | 47.5           | 45.8                    | 49.1  |
| 0.80                 | 49.7           | 48.0                    | 51.5  |
| 0.85                 | 52.5           | 50.5                    | 54.4  |
| 0.90                 | 56.1           | 53.8                    | 58.4  |
| 0.95                 | 62.0           | 59.1                    | 65.0  |
| 0.99                 | 75.0           | 70.4                    | 79.6  |
| Males                |                |                         |       |
| 0.05                 | 19.8           | 15.8                    | 23.7  |
| 0.10                 | 25.1           | 21.9                    | 28.4  |
| 0.15                 | 28.5           | 25.6                    | 31.3  |
| 0.20                 | 31.0           | 28.4                    | 33.5  |
| 0.25                 | 33.0           | 30.7                    | 35.4  |
| 0.30                 | 34.9           | 32.7                    | 37.0  |
| 0.35                 | 36.5           | 34.5                    | 38.5  |
| 0.40                 | 38.0           | 36.2                    | 39.9  |
| 0.45                 | 39.5           | 37.8                    | 41.2  |
| 0.50                 | 41.0           | 39.3                    | 42.6  |
| 0.55                 | 42.4           | 40.8                    | 44.0  |
| 0.60                 | 43.9           | 42.4                    | 45.4  |
| 0.65                 | 45.4           | 43.9                    | 46.9  |
| 0.70                 | 47.1           | 45.6                    | 48.5  |
| 0.75                 | 48.9           | 47.3                    | 50.4  |
| 0.80                 | 50.9           | 49.3                    | 52.6  |
| 0.85                 | 53.4           | 51.7                    | 55.2  |
| 0.90                 | 56.8           | 54.7                    | 58.9  |
| 0.95                 | 62.2           | 59.5                    | 64.8  |
| 0.99                 | 74.0           | 69.9                    | 78.2  |

<sup>a</sup> Offshore strata 13–25.

spring of 1985–1990, was 38.8 cm and 41.0 cm, respectively (Tables 3 and 5, Fig. 3). From the Gulf of Maine region,  $L_{50}$  values were 32.1 cm for females and 36.0 cm for males for Atlantic cod sampled during 1985–1990 (Tables 3 and 7, Fig. 5). Median age at maturity was 1.7 yr and 1.9 yr, respectively, for females and males from Georges Bank (Table 4, Fig. 4) and 2.1 yr and 2.3 yr for females and males from the Gulf of Maine (Table 4, Fig. 6); proportions of mature fish calculated for each age are presented in Tables 6 and 8.

Median maturity values from the present study are noticeably lower for both sexes compared with previous studies. Livingstone and Dery (1976) noted that  $L_{50}$  values for Atlantic cod collected from Georges Bank during the spring of 1972 were 51.5 cm and 44.0 cm for females and males, respectively. The  $A_{50}$  values were 2.9 yr and 2.6 yr for females and males, respectively. Atlan-

tic cod sampled over the entire region during 1977 (Morse 1979)<sup>2</sup> had  $L_{50}$  values of 49.6 cm and 53.7 cm for females and males, respectively.

Beacham (1983a) reported that slower-growing Atlantic cod of the Scotian Shelf area had  $L_{50}$  values of 52 cm for females and 51 cm for males in 1963 and that by 1978 the  $L_{50}$  values had declined to 35 cm and 38 cm, respectively. Median age values also declined from 3.7 yr and 4.8 yr for females and males, respectively, for the years 1959–1964 to 2.9 yr for females and 2.8 yrs for males during the years 1975–1979.

<sup>2</sup> W.W. Morse. 1979. An analysis of maturity observations of 12 groundfish species collected from Cape Hatteras, North Carolina to Nova Scotia in 1977. NMFS/NEFC Sandy Hook Laboratory Rep. No. SHL 79–32, 21 p.




**Figure 4**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of Georges Bank female ( $n=970$ ) and male ( $n=1021$ ) Atlantic cod, *Gadus morhua*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 6**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Georges Bank female ( $n=970$ ) and male ( $n=1021$ ) Atlantic cod, *Gadus morhua*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.23                 | 1.0  | 0.8                     | 1.2   |
| 0.60                 | 2.0  | 1.9                     | 2.1   |
| 0.88                 | 3.0  | 2.8                     | 3.2   |
| 0.97                 | 4.0  | 3.7                     | 4.3   |
| 0.99                 | 5.0  | 4.5                     | 5.5   |
| 1.00                 | 6.0  | 5.4                     | 6.6   |
| 1.00                 | 7.0  | 6.2                     | 7.8   |
| 1.00                 | 8.0  | 7.1                     | 8.9   |
| 1.00                 | 9.0  | 7.9                     | 10.1  |
| 1.00                 | 10.0 | 8.8                     | 11.2  |
| 1.00                 | 11.0 | 9.6                     | 12.4  |
| Males                |      |                         |       |
| 0.18                 | 1.0  | 0.8                     | 1.2   |
| 0.55                 | 2.0  | 1.9                     | 2.1   |
| 0.87                 | 3.0  | 2.8                     | 3.2   |
| 0.97                 | 4.0  | 3.7                     | 4.3   |
| 1.00                 | 5.0  | 4.6                     | 5.4   |
| 1.00                 | 6.0  | 5.4                     | 6.6   |
| 1.00                 | 7.0  | 6.3                     | 7.7   |
| 1.00                 | 8.0  | 7.1                     | 8.9   |
| 1.00                 | 9.0  | 8.0                     | 10.0  |
| 1.00                 | 10.0 | 8.9                     | 11.1  |
| 1.00                 | 11.0 | 9.7                     | 12.3  |
| 1.00                 | 12.0 | 10.6                    | 13.4  |

<sup>a</sup> Offshore strata 13–25.




**Figure 5**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of Gulf of Maine female ( $n=411$ ) and male ( $n=394$ ) Atlantic cod, *Gadus morhua*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 7**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Gulf of Maine female ( $n=411$ ) and male ( $n=394$ ) Atlantic cod, *Gadus morhua*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 14.9           | 10.4                    | 19.5  |
| 0.10                 | 19.3           | 15.6                    | 23.0  |
| 0.15                 | 22.0           | 18.8                    | 25.2  |
| 0.20                 | 24.0           | 21.2                    | 26.9  |
| 0.25                 | 25.7           | 23.1                    | 28.3  |
| 0.30                 | 27.2           | 24.8                    | 29.6  |
| 0.35                 | 28.5           | 26.3                    | 30.8  |
| 0.40                 | 29.8           | 27.7                    | 31.9  |
| 0.45                 | 31.0           | 29.0                    | 33.0  |
| 0.50                 | 32.1           | 30.2                    | 34.1  |
| 0.55                 | 33.3           | 31.4                    | 35.2  |
| 0.60                 | 34.5           | 32.6                    | 36.4  |
| 0.65                 | 35.8           | 33.9                    | 37.7  |
| 0.70                 | 37.1           | 35.1                    | 39.1  |
| 0.75                 | 38.6           | 36.5                    | 40.7  |
| 0.80                 | 40.3           | 38.0                    | 42.5  |
| 0.85                 | 42.3           | 39.7                    | 44.8  |
| 0.90                 | 45.0           | 42.0                    | 48.0  |
| 0.95                 | 49.4           | 45.6                    | 53.1  |
| 0.99                 | 59.0           | 53.4                    | 64.6  |
| Males                |                |                         |       |
| 0.05                 | 15.0           | 10.1                    | 19.9  |
| 0.10                 | 20.3           | 16.4                    | 24.2  |
| 0.15                 | 23.6           | 20.2                    | 27.0  |
| 0.20                 | 26.1           | 23.1                    | 29.1  |
| 0.25                 | 28.1           | 25.4                    | 30.8  |
| 0.30                 | 29.9           | 27.5                    | 32.4  |
| 0.35                 | 31.6           | 29.3                    | 33.8  |
| 0.40                 | 33.1           | 30.9                    | 35.2  |
| 0.45                 | 34.5           | 32.5                    | 36.6  |
| 0.50                 | 36.0           | 33.9                    | 38.0  |
| 0.55                 | 37.4           | 35.4                    | 39.4  |
| 0.60                 | 38.9           | 36.8                    | 40.9  |
| 0.65                 | 40.4           | 38.2                    | 42.5  |
| 0.70                 | 42.0           | 39.7                    | 44.3  |
| 0.75                 | 43.8           | 41.3                    | 46.2  |
| 0.80                 | 45.8           | 43.1                    | 48.6  |
| 0.85                 | 48.3           | 45.2                    | 51.4  |
| 0.90                 | 51.6           | 48.0                    | 55.2  |
| 0.95                 | 56.9           | 52.4                    | 61.5  |
| 0.99                 | 68.7           | 61.9                    | 75.5  |

<sup>a</sup> Offshore strata 26–30, 36–40.




**Figure 6**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of Gulf of Maine female ( $n=409$ ) and male ( $n=392$ ) Atlantic cod, *Gadus morhua*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 8**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Gulf of Maine female ( $n=409$ ) and male ( $n=392$ ) Atlantic cod, *Gadus morhua*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.08                 | 1.0  | 0.7                     | 1.3   |
| 0.44                 | 2.0  | 1.8                     | 2.2   |
| 0.88                 | 3.0  | 2.8                     | 3.2   |
| 0.99                 | 4.0  | 3.6                     | 4.4   |
| 1.00                 | 5.0  | 4.4                     | 5.6   |
| 1.00                 | 6.0  | 5.2                     | 6.8   |
| 1.00                 | 7.0  | 6.0                     | 8.0   |
| 1.00                 | 8.0  | 6.8                     | 9.2   |
| 1.00                 | 9.0  | 7.6                     | 10.4  |
| 1.00                 | 10.0 | 8.4                     | 11.6  |
| 1.00                 | 11.0 | 9.2                     | 12.8  |
| 1.00                 | 12.0 | 10.0                    | 14.0  |
| 1.00                 | 13.0 | 10.8                    | 15.2  |
| 1.00                 | 14.0 | 11.6                    | 16.4  |
| Males                |      |                         |       |
| 0.10                 | 1.0  | 0.7                     | 1.3   |
| 0.37                 | 2.0  | 1.8                     | 2.2   |
| 0.76                 | 3.0  | 2.8                     | 3.2   |
| 0.94                 | 4.0  | 3.6                     | 4.4   |
| 0.99                 | 5.0  | 4.5                     | 5.5   |
| 1.00                 | 6.0  | 5.3                     | 6.7   |
| 1.00                 | 7.0  | 6.1                     | 7.9   |
| 1.00                 | 8.0  | 6.9                     | 9.1   |
| 1.00                 | 9.0  | 7.6                     | 10.4  |

<sup>a</sup> Offshore strata 26–30, 36–40.

## Haddock: *Melanogrammus aeglefinus*

Haddock range from West Greenland to Cape Hatteras in the northwest Atlantic, but are most common on the Scotian Shelf and on Georges Bank. Two stocks exist off New England: the major spawning concentration occurs on Georges Bank and a second stock with more dispersed spawning occurs in the Gulf of Maine (Grosslein 1962). Spawning takes place during late winter and early spring with peak activity occurring in March and April (Table 2). Commercial landings of haddock from Georges Bank increased from a long-term annual average of 50,000 metric tons prior to the 1960's to about 150,000 t per year in 1965 and 1966, but declined precipitously thereafter to about 5,000 t annually during the early-to-mid-1970's; after a temporary increase to about 25,000 t per year during the late 1970's and early 1980's, total landings from this stock declined again to less than 10,000 t per year since 1985. Landings from the Gulf of Maine have generally remained below 5,000 t per year except during the early 1980's when the total ranged between 6,000 and 7,000 t

annually; landings have since declined to less than 1,000 t per year (NEFSC 1991).

Median length at maturity for female and male had-




**Figure 7**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of Georges Bank female ( $n=580$ ) and male ( $n=478$ ) haddock, *Melanogrammus aeglefinus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1989.

**Table 9**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Georges Bank female ( $n=580$ ) and male ( $n=478$ ) haddock, *Melanogrammus aeglefinus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 15.7           | 11.7                    | 19.7  |
| 0.10                 | 19.3           | 15.9                    | 22.6  |
| 0.15                 | 21.5           | 18.5                    | 24.4  |
| 0.20                 | 23.1           | 20.5                    | 25.8  |
| 0.25                 | 24.5           | 22.1                    | 26.9  |
| 0.30                 | 25.7           | 23.5                    | 27.9  |
| 0.35                 | 26.8           | 24.7                    | 28.8  |
| 0.40                 | 27.8           | 25.9                    | 29.7  |
| 0.45                 | 28.8           | 27.0                    | 30.6  |
| 0.50                 | 29.7           | 28.0                    | 31.4  |
| 0.55                 | 30.7           | 29.1                    | 32.3  |
| 0.60                 | 31.6           | 30.1                    | 33.2  |
| 0.65                 | 32.7           | 31.2                    | 34.1  |
| 0.70                 | 33.7           | 32.3                    | 35.2  |
| 0.75                 | 34.9           | 33.5                    | 36.4  |
| 0.80                 | 36.3           | 34.8                    | 37.8  |
| 0.85                 | 38.0           | 36.3                    | 39.6  |
| 0.90                 | 40.2           | 38.2                    | 42.1  |
| 0.95                 | 43.7           | 41.2                    | 46.2  |
| 0.99                 | 51.6           | 47.7                    | 55.5  |
| Males                |                |                         |       |
| 0.05                 | 11.2           | 6.0                     | 16.4  |
| 0.10                 | 15.2           | 10.9                    | 19.5  |
| 0.15                 | 17.6           | 13.8                    | 21.4  |
| 0.20                 | 19.5           | 16.0                    | 22.9  |
| 0.25                 | 21.0           | 17.9                    | 24.1  |
| 0.30                 | 22.3           | 19.4                    | 25.2  |
| 0.35                 | 23.5           | 20.9                    | 26.2  |
| 0.40                 | 24.7           | 22.2                    | 27.1  |
| 0.45                 | 25.8           | 23.5                    | 28.1  |
| 0.50                 | 26.8           | 24.7                    | 29.0  |
| 0.55                 | 27.9           | 25.9                    | 29.9  |
| 0.60                 | 29.0           | 27.1                    | 30.9  |
| 0.65                 | 30.1           | 28.3                    | 31.9  |
| 0.70                 | 31.3           | 29.5                    | 33.1  |
| 0.75                 | 32.6           | 30.9                    | 34.4  |
| 0.80                 | 34.2           | 32.4                    | 36.0  |
| 0.85                 | 36.0           | 34.0                    | 38.0  |
| 0.90                 | 38.5           | 36.2                    | 40.7  |
| 0.95                 | 42.4           | 39.5                    | 45.4  |
| 0.99                 | 51.2           | 46.4                    | 55.9  |

<sup>a</sup> Offshore strata 13–25, 29–30.

dock from Georges Bank was 29.7 and 26.8 cm, respectively (Tables 3 and 9, Fig. 7). For haddock from the Gulf of Maine region, values of  $L_{50}$  were 34.5 cm for females and 35.0 cm for males (Tables 3 and 11, Fig. 9). Median age at maturity was 1.5 yr for females and 1.3 yr for males from the former region, and 1.8 yr for females and 2.1 yr for males from the latter region (Table 4, Figs. 8 and 10); proportions of mature fish calculated for each age are presented in Tables 10 and 12.

Values of  $L_{50}$  above are considerably lower for both sexes than those reported by Morse (1979)<sup>2</sup> for the year 1977 over the entire region. Overholtz (1987), however, concluded that  $L_{50}$  values for both sexes on

Georges Bank had increased between 1977 and 1983, a period dominated by the relatively large 1975 and 1978 year classes. For the overall 1977–1983 period, the median length at maturity on Georges Bank was 40 cm for females and 37 cm for males. On the southwestern Scotian Shelf, however, median age at maturity declined substantially between the early 1960's and late 1970's from 4.6 to 3.0 yr for females and from 4.7 to 2.8 yr for males (Beacham 1983b). Templeman et al. (1978) and Templeman and Bishop (1979) also noted substantial changes in both  $L_{50}$  and  $A_{50}$  for haddock on the Grand Bank and St. Pierre Bank, and attributed these to interactions between growth and exploitation.




**Figure 8**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of Georges Bank female ( $n=580$ ) and male ( $n=478$ ) haddock, *Melanogrammus aeglefinus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1989.

**Table 10**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Georges Bank female ( $n=580$ ) and male ( $n=478$ ) haddock, *Melanogrammus aeglefinus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.26                 | 1.0  | 0.8                     | 1.2   |
| 0.77                 | 2.0  | 1.9                     | 2.1   |
| 0.97                 | 3.0  | 2.7                     | 3.3   |
| 1.00                 | 4.0  | 3.5                     | 4.5   |
| 1.00                 | 5.0  | 4.3                     | 5.7   |
| 1.00                 | 6.0  | 5.1                     | 6.9   |
| 1.00                 | 7.0  | 5.9                     | 8.1   |
| 1.00                 | 8.0  | 6.6                     | 9.4   |
| 1.00                 | 9.0  | 7.4                     | 10.6  |
| 1.00                 | 10.0 | 8.2                     | 11.8  |
| Males                |      |                         |       |
| 0.36                 | 1.0  | 0.8                     | 1.2   |
| 0.83                 | 2.0  | 1.8                     | 2.2   |
| 0.98                 | 3.0  | 2.6                     | 3.4   |
| 1.00                 | 4.0  | 3.4                     | 4.6   |
| 1.00                 | 5.0  | 4.2                     | 5.8   |
| 1.00                 | 6.0  | 4.9                     | 7.1   |
| 1.00                 | 7.0  | 5.7                     | 8.3   |
| 1.00                 | 8.0  | 6.4                     | 9.6   |
| 1.00                 | 9.0  | 7.2                     | 10.8  |
| 1.00                 | 10.0 | 7.9                     | 12.1  |

<sup>a</sup> Offshore strata 13–25, 29–30.




**Figure 9**

Calculated and observed (•) proportion mature at length based on macroscopic observations of whole gonads of Gulf of Maine female ( $n=69$ ) and male ( $n=49$ ) haddock, *Melanogrammus aeglefinus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1989.

**Table 11**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Gulf of Maine female ( $n=69$ ) and male ( $n=49$ ) haddock, *Melanogrammus aeglefinus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 29.0           | 18.8                    | 39.2  |
| 0.10                 | 30.4           | 21.8                    | 39.1  |
| 0.15                 | 31.3           | 23.5                    | 39.0  |
| 0.20                 | 31.9           | 24.9                    | 39.0  |
| 0.25                 | 32.5           | 25.9                    | 39.0  |
| 0.30                 | 32.9           | 26.9                    | 39.0  |
| 0.35                 | 33.3           | 27.7                    | 39.0  |
| 0.40                 | 33.7           | 28.5                    | 39.0  |
| 0.45                 | 34.1           | 29.2                    | 39.0  |
| 0.50                 | 34.5           | 29.9                    | 39.1  |
| 0.55                 | 34.9           | 30.6                    | 39.1  |
| 0.60                 | 35.2           | 31.3                    | 39.2  |
| 0.65                 | 35.6           | 32.0                    | 39.3  |
| 0.70                 | 36.1           | 32.6                    | 39.5  |
| 0.75                 | 36.5           | 33.3                    | 39.7  |
| 0.80                 | 37.1           | 34.0                    | 40.1  |
| 0.85                 | 37.7           | 34.6                    | 40.8  |
| 0.90                 | 38.6           | 35.2                    | 41.9  |
| 0.95                 | 39.9           | 35.7                    | 44.2  |
| 0.99                 | 43.0           | 35.9                    | 50.2  |
| Males                |                |                         |       |
| 0.05                 | 27.2           | 18.8                    | 35.5  |
| 0.10                 | 29.2           | 22.2                    | 36.1  |
| 0.15                 | 30.4           | 24.3                    | 36.5  |
| 0.20                 | 31.3           | 25.9                    | 36.8  |
| 0.25                 | 32.1           | 27.1                    | 37.0  |
| 0.30                 | 32.8           | 28.2                    | 37.3  |
| 0.35                 | 33.4           | 29.2                    | 37.6  |
| 0.40                 | 33.9           | 30.0                    | 37.9  |
| 0.45                 | 34.5           | 30.8                    | 38.1  |
| 0.50                 | 35.0           | 31.6                    | 38.5  |
| 0.55                 | 35.6           | 32.3                    | 38.8  |
| 0.60                 | 36.1           | 33.0                    | 39.2  |
| 0.65                 | 36.7           | 33.6                    | 39.7  |
| 0.70                 | 37.3           | 34.3                    | 40.3  |
| 0.75                 | 38.0           | 34.9                    | 41.0  |
| 0.80                 | 38.7           | 35.5                    | 42.0  |
| 0.85                 | 39.7           | 36.0                    | 43.3  |
| 0.90                 | 40.9           | 36.7                    | 45.1  |
| 0.95                 | 42.9           | 37.5                    | 48.3  |
| 0.99                 | 47.3           | 38.8                    | 55.8  |

<sup>a</sup> Offshore strata 26–28, 36–40.




**Figure 10**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of Gulf of Maine female ( $n=69$ ) and male ( $n=49$ ) haddock, *Melanogrammus aeglefinus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1989.

**Table 12**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Gulf of Maine female ( $n=69$ ) and male ( $n=49$ ) haddock, *Melanogrammus aeglefinus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.13                 | 1.0  | 0.0                     | 2.0   |
| 0.64                 | 2.0  | 1.4                     | 2.6   |
| 0.96                 | 3.0  | 2.4                     | 3.6   |
| 1.00                 | 4.0  | 2.9                     | 5.1   |
| 1.00                 | 5.0  | 3.3                     | 6.7   |
| 1.00                 | 6.0  | 3.7                     | 8.3   |
| 1.00                 | 7.0  | 4.0                     | 10.0  |
| 1.00                 | 8.0  | 4.4                     | 11.6  |
| 1.00                 | 9.0  | 4.8                     | 13.2  |
| 1.00                 | 10.0 | 5.2                     | 14.8  |
| Males                |      |                         |       |
| 0.10                 | 1.0  | -0.3                    | 2.3   |
| 0.43                 | 2.0  | 1.3                     | 2.7   |
| 0.85                 | 3.0  | 2.5                     | 3.5   |
| 0.97                 | 4.0  | 2.9                     | 5.1   |
| 1.00                 | 5.0  | 3.3                     | 6.7   |
| 1.00                 | 6.0  | 3.6                     | 8.4   |
| 1.00                 | 7.0  | 3.9                     | 10.1  |

<sup>a</sup> Offshore strata 26–28, 36–40.

## Pollock: *Pollachius virens*

Pollock occur on both sides of the north Atlantic. In the northwest Atlantic they are most abundant on the Scotian Shelf and in the Gulf of Maine (Mayo et al. 1989). Spawning takes place during winter with peak activity occurring in December and January (Table 2). Total landings of pollock from the Scotian Shelf-Gulf of Maine region increased sharply from about 38,000 metric tons per year in the mid 1970's to over 68,000 t in 1986, but have since declined to about 50,000 t in 1989 and 1990 (NEFSC 1991).

Median length at maturity for female and male pollock from the Gulf of Maine was 39.1 and 41.8 cm, respectively (Tables 3 and 13, Fig. 11). Median age at maturity was 2.0 yr for females and 2.3 yr for males (Table 4, Fig. 12); proportions of mature fish calculated for each age are presented in Table 14.

Values of  $L_{50}$  and  $A_{50}$  above are lower than those of Mayo et al. (1989) who reported  $L_{50}$  estimates of 47-48 cm for females and 50.5 cm for males, based on USA and Canadian samples collected between 1977 and 1984.




**Figure 11**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=154$ ) and male ( $n=143$ ) pollock, *Pollachius virens*, sampled during Northeast Fisheries Science Center autumn bottom trawl surveys, 1986–1988.

Estimates of  $A_{50}$  reported by Mayo et al. (1989) (3.2 yr for females and 3.5 yr for males) are at least one full year higher than those given above. Steele (1963) re-

**Table 13**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=154$ ) and male ( $n=143$ ) pollock, *Pollachius virens*, sampled during Northeast Fisheries Science Center autumn bottom trawl surveys<sup>a</sup>, 1986–1988.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 19.9           | 12.0                    | 27.8  |
| 0.10                 | 24.8           | 18.4                    | 31.2  |
| 0.15                 | 27.8           | 22.2                    | 33.4  |
| 0.20                 | 30.1           | 25.1                    | 35.0  |
| 0.25                 | 31.9           | 27.4                    | 36.4  |
| 0.30                 | 33.6           | 29.4                    | 37.7  |
| 0.35                 | 35.1           | 31.2                    | 38.9  |
| 0.40                 | 36.5           | 32.8                    | 40.1  |
| 0.45                 | 37.8           | 34.3                    | 41.3  |
| 0.50                 | 39.1           | 35.7                    | 42.5  |
| 0.55                 | 40.4           | 37.1                    | 43.7  |
| 0.60                 | 41.7           | 38.4                    | 45.1  |
| 0.65                 | 43.1           | 39.7                    | 46.5  |
| 0.70                 | 44.6           | 41.0                    | 48.2  |
| 0.75                 | 46.3           | 42.4                    | 50.1  |
| 0.80                 | 48.1           | 44.0                    | 52.3  |
| 0.85                 | 50.4           | 45.7                    | 55.1  |
| 0.90                 | 53.4           | 47.9                    | 58.9  |
| 0.95                 | 58.3           | 51.4                    | 65.2  |
| 0.99                 | 69.0           | 58.7                    | 79.3  |
| Males                |                |                         |       |
| 0.05                 | 23.7           | 17.1                    | 30.2  |
| 0.10                 | 28.3           | 23.0                    | 33.5  |
| 0.15                 | 31.1           | 26.6                    | 35.6  |
| 0.20                 | 33.3           | 29.2                    | 37.3  |
| 0.25                 | 35.0           | 31.3                    | 38.7  |
| 0.30                 | 36.6           | 33.1                    | 40.0  |
| 0.35                 | 38.0           | 34.7                    | 41.2  |
| 0.40                 | 39.3           | 36.1                    | 42.4  |
| 0.45                 | 40.5           | 37.4                    | 43.7  |
| 0.50                 | 41.8           | 38.7                    | 44.9  |
| 0.55                 | 43.0           | 39.8                    | 46.2  |
| 0.60                 | 44.3           | 41.0                    | 47.6  |
| 0.65                 | 45.6           | 42.1                    | 49.1  |
| 0.70                 | 47.0           | 43.3                    | 50.7  |
| 0.75                 | 48.5           | 44.5                    | 52.6  |
| 0.80                 | 50.3           | 45.9                    | 54.8  |
| 0.85                 | 52.5           | 47.5                    | 57.4  |
| 0.90                 | 55.3           | 49.5                    | 61.1  |
| 0.95                 | 59.9           | 52.8                    | 67.0  |
| 0.99                 | 70.1           | 59.8                    | 80.4  |

<sup>a</sup> Offshore strata 13–34, 351, 36–40.

ported median maturation values of 62.5 cm at 5–6 yrs for females and 58 cm at 4–5 yrs for males collected from the Bay of Fundy in 1960 and 1961. The corresponding median ages are similar to those presented by Beacham (1983c) for a similar time period from the

southern Scotian Shelf. Beacham further noted substantial declines in pollock  $L_{50}$  and  $A_{50}$  values of 5–6 cm and 1–2 yr between the early 1960's and late 1970's in the same region.




**Figure 12**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=154$ ) and male ( $n=143$ ) pollock, *Pollachius virens*, sampled during Northeast Fisheries Science Center autumn bottom trawl surveys, 1986–1988.

**Table 14**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=154$ ) and male ( $n=143$ ) pollock, *Pollachius virens*, sampled during Northeast Fisheries Science Center autumn bottom trawl surveys<sup>a</sup>, 1986–1988.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.24                 | 1.0  | 0.4                     | 1.6   |
| 0.50                 | 2.0  | 1.6                     | 2.4   |
| 0.76                 | 3.0  | 2.5                     | 3.5   |
| 0.91                 | 4.0  | 3.3                     | 4.7   |
| 0.97                 | 5.0  | 3.9                     | 6.1   |
| 0.99                 | 6.0  | 4.6                     | 7.4   |
| 1.00                 | 7.0  | 5.2                     | 8.8   |
| 1.00                 | 8.0  | 5.8                     | 10.2  |
| 1.00                 | 9.0  | 6.5                     | 11.5  |
| 1.00                 | 10.0 | 7.1                     | 12.9  |
| 1.00                 | 11.0 | 7.7                     | 14.3  |
| 1.00                 | 12.0 | 8.4                     | 15.6  |
| 1.00                 | 13.0 | 9.0                     | 17.0  |
| 1.00                 | 14.0 | 9.6                     | 18.4  |
| 1.00                 | 15.0 | 10.2                    | 19.8  |
| Males                |      |                         |       |
| 0.07                 | 1.0  | 0.5                     | 1.5   |
| 0.35                 | 2.0  | 1.7                     | 2.3   |
| 0.78                 | 3.0  | 2.6                     | 3.4   |
| 0.96                 | 4.0  | 3.3                     | 4.7   |
| 0.99                 | 5.0  | 4.0                     | 6.0   |
| 1.00                 | 6.0  | 4.6                     | 7.4   |
| 1.00                 | 7.0  | 5.3                     | 8.7   |
| 1.00                 | 8.0  | 5.9                     | 10.1  |
| 1.00                 | 9.0  | 6.5                     | 11.5  |
| 1.00                 | 10.0 | 7.2                     | 12.8  |
| 1.00                 | 11.0 | 7.8                     | 14.2  |
| 1.00                 | 12.0 | 8.5                     | 15.5  |
| 1.00                 | 13.0 | 9.1                     | 16.9  |
| 1.00                 | 14.0 | 9.7                     | 18.3  |
| 1.00                 | 15.0 | 10.4                    | 19.6  |
| 1.00                 | 16.0 | 11.0                    | 21.0  |
| 1.00                 | 17.0 | 11.7                    | 22.3  |
| 1.00                 | 18.0 | 12.3                    | 23.7  |

<sup>a</sup> Offshore strata 13–34, 351, 36–40.

## White Hake: *Urophycis tenuis*

White hake are widely distributed from Labrador to Georges Bank, and in deeper slope waters southward to Florida. In U.S. waters, white hake of the Gulf of Maine-Georges Bank region are considered to be a unit stock. Information relative to white hake spawning in this region is both limited and conflicting, although ripe fish have been observed from October to May (Table 2). Commercial landings of white hake averaged 4,700 metric tons per year during 1975–1980, increased to an average of 6,700 t from 1981 to 1988, and subsequently declined to 5,600 t in 1989 (NEFSC 1991).

Median length at maturity for female and male white hake was 35.1 and 32.7 cm, respectively (Tables 3 and 15, Fig. 13). Median age at maturity was 1.4 yr for both sexes (Table 4, Fig. 14); proportions of mature fish calculated for each age are presented in Table 16.

Values of  $L_{50}$  above are considerably lower than those obtained by Morse (1979)<sup>2</sup> from 1977 data (47.6 and 43.0 cm for females and males, respectively) and by Burnett et al. (1984)<sup>3</sup> from data collected during 1977–

<sup>3</sup> J. Burnett, S.H. Clark, and L. O'Brien. 1984. A preliminary assessment of white hake in the Gulf of Maine-Georges Bank area. NMFS/NEFC, Woods Hole Lab. Ref. Doc. No. 84–31, 33 p.

**Table 15**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=455$ ) and male ( $n=346$ ) white hake, *Urophycis tenuis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1987–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 13.6           | 8.4                     | 18.8  |
| 0.10                 | 19.1           | 14.9                    | 23.3  |
| 0.15                 | 22.5           | 18.8                    | 26.1  |
| 0.20                 | 25.0           | 21.8                    | 28.3  |
| 0.25                 | 27.1           | 24.2                    | 30.0  |
| 0.30                 | 29.0           | 26.3                    | 31.6  |
| 0.35                 | 30.6           | 28.2                    | 33.1  |
| 0.40                 | 32.2           | 29.9                    | 34.5  |
| 0.45                 | 33.7           | 31.6                    | 35.8  |
| 0.50                 | 35.1           | 33.1                    | 37.2  |
| 0.55                 | 36.6           | 34.7                    | 38.6  |
| 0.60                 | 38.1           | 36.2                    | 40.0  |
| 0.65                 | 39.7           | 37.7                    | 41.6  |
| 0.70                 | 41.3           | 39.3                    | 43.4  |
| 0.75                 | 43.2           | 41.0                    | 45.3  |
| 0.80                 | 45.3           | 42.9                    | 47.7  |
| 0.85                 | 47.8           | 45.1                    | 50.5  |
| 0.90                 | 51.2           | 48.0                    | 54.4  |
| 0.95                 | 56.7           | 52.6                    | 60.8  |
| 0.99                 | 68.7           | 62.5                    | 75.0  |
| Males                |                |                         |       |
| 0.05                 | 20.3           | 17.1                    | 23.6  |
| 0.10                 | 23.5           | 20.9                    | 26.1  |
| 0.15                 | 25.4           | 23.2                    | 27.7  |
| 0.20                 | 26.9           | 24.9                    | 28.9  |
| 0.25                 | 28.1           | 26.3                    | 29.9  |
| 0.30                 | 29.2           | 27.5                    | 30.8  |
| 0.35                 | 30.1           | 28.6                    | 31.6  |
| 0.40                 | 31.0           | 29.6                    | 32.4  |
| 0.45                 | 31.9           | 30.5                    | 33.2  |
| 0.50                 | 32.7           | 31.4                    | 34.0  |
| 0.55                 | 33.6           | 32.3                    | 34.8  |
| 0.60                 | 34.4           | 33.2                    | 35.6  |
| 0.65                 | 35.3           | 34.1                    | 36.5  |
| 0.70                 | 36.3           | 35.0                    | 37.5  |
| 0.75                 | 37.3           | 36.0                    | 38.7  |
| 0.80                 | 38.5           | 37.0                    | 40.0  |
| 0.85                 | 40.0           | 38.3                    | 41.7  |
| 0.90                 | 41.9           | 39.9                    | 44.0  |
| 0.95                 | 45.1           | 42.5                    | 47.7  |
| 0.99                 | 52.0           | 48.1                    | 55.9  |

<sup>a</sup> Offshore strata 21–30, 33–34, 351, 36–40.




**Figure 13**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=455$ ) and male ( $n=346$ ) white hake, *Urophycis tenuis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1987–1989.

82 (42 cm for both sexes) for the Gulf of Maine-Georges Bank region. Musick (1969) noted that females matured at 48 to 64 cm and males at 40 to 52 cm, but his data included fish from both U.S. and Canadian waters.

Maturation of white hake in the Gulf of Maine-Georges Bank region occurs at a smaller size and younger age than for slower-growing white hake in Canadian waters. Battle (1951) observed no spawning females below 57 cm in size and 4 yr in age within the Bay of Fundy, and Beacham and Nepszy (1980) and Beacham (1983c) reported values of  $L_{50}$  ranging from 40.1 to 53.9 cm for females and from 36.6 to 44.1 cm for males in the Gulf of St. Lawrence and Scotian Shelf regions during 1971-78, corresponding to ages of 3 to 5 yr.




**Figure 14**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=455$ ) and male ( $n=346$ ) white hake, *Urophycis tenuis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1987-1989.

However, maturation of white hake in the Gulf of Maine-Georges Bank region may be a moot point in view of evidence that the species does not successfully spawn in that region. Musick (1969) observed that spawning was weak and aperiodic, and hypothesized that recruits from the Scotian Shelf sustained the population. More recently, Fahay and Able (1989) reported the absence of white hake eggs and larvae in the Gulf of Maine, and presented evidence which suggested that the local population was sustained primarily by recruits from a spring-spawning continental slope population and secondarily by recruits from the summer-spawning and fall-spawning Scotian Shelf and southern Gulf of St. Lawrence populations.

**Table 16**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=455$ ) and male ( $n=346$ ) white hake, *Urophycis tenuis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1987-1989.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.33                 | 1.0  | 0.8                     | 1.2   |
| 0.74                 | 2.0  | 1.8                     | 2.2   |
| 0.94                 | 3.0  | 2.7                     | 3.3   |
| 0.99                 | 4.0  | 3.5                     | 4.5   |
| 1.00                 | 5.0  | 4.3                     | 5.7   |
| 1.00                 | 6.0  | 5.1                     | 6.9   |
| 1.00                 | 7.0  | 5.9                     | 8.1   |
| 1.00                 | 8.0  | 6.7                     | 9.3   |
| 1.00                 | 9.0  | 7.5                     | 10.5  |
| 1.00                 | 10.0 | 8.3                     | 11.7  |
| 1.00                 | 11.0 | 9.1                     | 12.9  |
| 1.00                 | 12.0 | 9.9                     | 14.1  |
| 1.00                 | 13.0 | 10.7                    | 15.3  |
| 1.00                 | 14.0 | 11.5                    | 16.5  |
| 1.00                 | 15.0 | 12.3                    | 17.7  |
| 1.00                 | 16.0 | 13.1                    | 18.9  |
| 1.00                 | 17.0 | 13.9                    | 20.1  |
| 1.00                 | 18.0 | 14.7                    | 21.3  |
| Males                |      |                         |       |
| 0.29                 | 1.0  | 0.8                     | 1.2   |
| 0.82                 | 2.0  | 1.8                     | 2.2   |
| 0.98                 | 3.0  | 2.7                     | 3.3   |
| 1.00                 | 4.0  | 3.4                     | 4.6   |
| 1.00                 | 5.0  | 4.2                     | 5.8   |
| 1.00                 | 6.0  | 5.0                     | 7.0   |
| 1.00                 | 7.0  | 5.8                     | 8.2   |

<sup>a</sup> Offshore strata 21-30, 33-34, 351, 36-40.

## Red Hake: *Urophycis chuss*

Red hake are distributed from the Gulf of St. Lawrence to Cape Hatteras, North Carolina. It is assumed that two stocks exist in U.S. waters, one occupying the Gulf of Maine and the northern portion of Georges Bank, and the other situated on southern Georges Bank and throughout the Southern New England-Middle Atlantic region. Spawning occurs from May to September for the northern stock, with peak activity during June and July, while the southern stock spawns over a protracted period from March to October (Table 2). Commercial landings for this under-exploited species have averaged 1,000 metric tons per year since 1980 for the northern stock, whereas landings from the southern stock have decreased from over 4,000 t in 1980 to 800 t in 1989 (NEFSC 1991).

Median length at maturity for female and male red hake from the Gulf of Maine-northern Georges Bank region was 26.9 and 22.2 cm, respectively (Tables 3 and 17, Fig. 15). For red hake from the Mid-Atlantic-south-

ern Georges Bank region, values of  $L_{50}$  were 25.1 cm for females and 23.8 cm for males (Tables 3 and 19, Fig.

**Table 17**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=667$ ) and male ( $n=595$ ) red hake, *Urophycis chuss*, from the Gulf of Maine-northern Georges Bank region sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1982–1985.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 21.0           | 19.2                    | 22.9  |
| 0.10                 | 22.5           | 21.0                    | 24.1  |
| 0.15                 | 23.4           | 22.1                    | 24.8  |
| 0.20                 | 24.1           | 22.9                    | 25.4  |
| 0.25                 | 24.7           | 23.5                    | 25.9  |
| 0.30                 | 25.2           | 24.1                    | 26.3  |
| 0.35                 | 25.7           | 24.6                    | 26.7  |
| 0.40                 | 26.1           | 25.1                    | 27.0  |
| 0.45                 | 26.5           | 25.6                    | 27.4  |
| 0.50                 | 26.9           | 26.0                    | 27.7  |
| 0.55                 | 27.3           | 26.5                    | 28.1  |
| 0.60                 | 27.7           | 26.9                    | 28.5  |
| 0.65                 | 28.1           | 27.3                    | 28.9  |
| 0.70                 | 28.6           | 27.8                    | 29.3  |
| 0.75                 | 29.1           | 28.3                    | 29.8  |
| 0.80                 | 29.6           | 28.8                    | 30.4  |
| 0.85                 | 30.3           | 29.5                    | 31.2  |
| 0.90                 | 31.2           | 30.3                    | 32.2  |
| 0.95                 | 32.7           | 31.5                    | 33.9  |
| 0.99                 | 36.0           | 34.2                    | 37.8  |
| Males                |                |                         |       |
| 0.05                 | 16.8           | 14.5                    | 19.1  |
| 0.10                 | 18.2           | 16.2                    | 20.1  |
| 0.15                 | 19.0           | 17.3                    | 20.8  |
| 0.20                 | 19.7           | 18.1                    | 21.3  |
| 0.25                 | 20.2           | 18.7                    | 21.7  |
| 0.30                 | 20.7           | 19.3                    | 22.1  |
| 0.35                 | 21.1           | 19.8                    | 22.4  |
| 0.40                 | 21.5           | 20.2                    | 22.7  |
| 0.45                 | 21.9           | 20.7                    | 23.0  |
| 0.50                 | 22.2           | 21.1                    | 23.3  |
| 0.55                 | 22.6           | 21.5                    | 23.7  |
| 0.60                 | 23.0           | 22.0                    | 24.0  |
| 0.65                 | 23.4           | 22.4                    | 24.3  |
| 0.70                 | 23.8           | 22.8                    | 24.7  |
| 0.75                 | 24.2           | 23.3                    | 25.2  |
| 0.80                 | 24.8           | 23.8                    | 25.7  |
| 0.85                 | 25.4           | 24.5                    | 26.4  |
| 0.90                 | 26.3           | 25.2                    | 27.3  |
| 0.95                 | 27.6           | 26.4                    | 28.9  |
| 0.99                 | 30.7           | 28.8                    | 32.6  |

<sup>a</sup> Offshore strata 20–30, 36–40.




**Figure 15**

Calculated and observed (•) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=667$ ) and male ( $n=595$ ) red hake, *Urophycis chuss*, from the Gulf of Maine-northern Georges Bank region sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1982–1985.

17). Median age at maturity was 1.8 yr for females and 1.4 yr for males in the former region and 1.7 yr for females and 1.8 yr for males in the latter region (Table 4, Figs. 16 and 18); proportions of mature fish calculated for each age are presented in Tables 18 and 20.

Morse (1979)<sup>2</sup> reported an  $L_{50}$  of 27.8 and 24.8 cm, respectively, for female and male red hake in the same

general region during 1977 but did not distinguish between stock areas. Musick (1969) observed that red hake matured at 2 yr of age, with females maturing at 27 to 33 cm and males at 24 to 30 cm. Markle et al. (1982) also reported that maturation occurred essentially at age 2, with most fish mature by 30 cm.




**Figure 16**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=667$ ) and male ( $n=595$ ) red hake, *Urophycis chuss*, from the Gulf of Maine-northern Georges Bank region sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1982–1985.

**Table 18**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=667$ ) and male ( $n=595$ ) red hake *Urophycis chuss* from the Gulf of Maine-northern Georges Bank region sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1982–1985.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.04                 | 1.0  | 0.7                     | 1.3   |
| 0.73                 | 2.0  | 1.9                     | 2.1   |
| 0.99                 | 3.0  | 2.7                     | 3.3   |
| 1.00                 | 4.0  | 3.4                     | 4.6   |
| 1.00                 | 5.0  | 4.2                     | 5.8   |
| 1.00                 | 6.0  | 4.9                     | 7.1   |
| 1.00                 | 7.0  | 5.6                     | 8.4   |
| 1.00                 | 8.0  | 6.3                     | 9.7   |
| 1.00                 | 9.0  | 7.1                     | 10.9  |
| 1.00                 | 10.0 | 7.8                     | 12.2  |
| 1.00                 | 11.0 | 8.5                     | 13.5  |
| 1.00                 | 12.0 | 9.2                     | 14.8  |
| Males                |      |                         |       |
| 0.15                 | 1.0  | 0.8                     | 1.2   |
| 0.91                 | 2.0  | 1.9                     | 2.1   |
| 1.00                 | 3.0  | 2.7                     | 3.3   |
| 1.00                 | 4.0  | 3.5                     | 4.5   |
| 1.00                 | 5.0  | 4.2                     | 5.8   |
| 1.00                 | 6.0  | 5.0                     | 7.0   |
| 1.00                 | 7.0  | 5.8                     | 8.2   |
| 1.00                 | 8.0  | 6.5                     | 9.5   |
| 1.00                 | 9.0  | 7.3                     | 10.7  |
| 1.00                 | 10.0 | 8.1                     | 11.9  |
| 1.00                 | 11.0 | 8.8                     | 13.2  |
| 1.00                 | 12.0 | 9.6                     | 14.4  |

<sup>a</sup> Offshore strata 20–30, 36–40.




**Figure 17**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=1020$ ) and male ( $n=753$ ) red hake, *Urophycis chuss*, from southern Georges Bank - Middle Atlantic sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1982–1985.

**Table 19**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=1020$ ) and male ( $n=753$ ) red hake, *Urophycis chuss*, from southern Georges Bank - Middle Atlantic sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1982–1985.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 19.3           | 17.9                    | 20.7  |
| 0.10                 | 20.8           | 19.6                    | 22.0  |
| 0.15                 | 21.7           | 20.6                    | 22.7  |
| 0.20                 | 22.4           | 21.4                    | 23.3  |
| 0.25                 | 22.9           | 22.1                    | 23.8  |
| 0.30                 | 23.4           | 22.6                    | 24.3  |
| 0.35                 | 23.9           | 23.1                    | 24.7  |
| 0.40                 | 24.3           | 23.6                    | 25.0  |
| 0.45                 | 24.7           | 24.0                    | 25.4  |
| 0.50                 | 25.1           | 24.4                    | 25.8  |
| 0.55                 | 25.5           | 24.8                    | 26.2  |
| 0.60                 | 25.9           | 25.3                    | 26.5  |
| 0.65                 | 26.3           | 25.7                    | 26.9  |
| 0.70                 | 26.8           | 26.2                    | 27.4  |
| 0.75                 | 27.3           | 26.6                    | 27.9  |
| 0.80                 | 27.8           | 27.2                    | 28.5  |
| 0.85                 | 28.5           | 27.8                    | 29.2  |
| 0.90                 | 29.4           | 28.7                    | 30.2  |
| 0.95                 | 30.9           | 30.0                    | 31.8  |
| 0.99                 | 34.1           | 32.8                    | 35.5  |
| Males                |                |                         |       |
| 0.05                 | 17.8           | 16.2                    | 19.3  |
| 0.10                 | 19.3           | 18.0                    | 20.6  |
| 0.15                 | 20.2           | 19.1                    | 21.4  |
| 0.20                 | 21.0           | 19.9                    | 22.0  |
| 0.25                 | 21.5           | 20.6                    | 22.5  |
| 0.30                 | 22.1           | 21.2                    | 23.0  |
| 0.35                 | 22.5           | 21.7                    | 23.4  |
| 0.40                 | 23.0           | 22.2                    | 23.7  |
| 0.45                 | 23.4           | 22.6                    | 24.1  |
| 0.50                 | 23.8           | 23.1                    | 24.5  |
| 0.55                 | 24.2           | 23.5                    | 24.9  |
| 0.60                 | 24.6           | 24.0                    | 25.3  |
| 0.65                 | 25.1           | 24.5                    | 25.7  |
| 0.70                 | 25.5           | 24.9                    | 26.1  |
| 0.75                 | 26.0           | 25.5                    | 26.6  |
| 0.80                 | 26.6           | 26.0                    | 27.2  |
| 0.85                 | 27.3           | 26.7                    | 28.0  |
| 0.90                 | 28.3           | 27.6                    | 29.0  |
| 0.95                 | 29.8           | 28.9                    | 30.7  |
| 0.99                 | 33.2           | 31.8                    | 34.6  |

<sup>a</sup> Inshore strata 1–46, 52, 55, offshore strata 1–19, 61–76.




**Figure 18**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=1020$ ) and male ( $n=753$ ) red hake, *Urophycis chuss*, from southern Georges Bank - Middle Atlantic sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1982–1985.

**Table 20**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=1020$ ) and male ( $n=753$ ) red hake, *Urophycis chuss*, from southern Georges Bank - Middle Atlantic sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1982–1985.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.09                 | 1.0  | 0.8                     | 1.2   |
| 0.70                 | 2.0  | 1.9                     | 2.1   |
| 0.98                 | 3.0  | 2.8                     | 3.2   |
| 1.00                 | 4.0  | 3.7                     | 4.3   |
| 1.00                 | 5.0  | 4.5                     | 5.5   |
| 1.00                 | 6.0  | 5.4                     | 6.6   |
| 1.00                 | 7.0  | 6.2                     | 7.8   |
| 1.00                 | 8.0  | 7.1                     | 8.9   |
| 1.00                 | 9.0  | 7.9                     | 10.1  |
| 1.00                 | 10.0 | 8.8                     | 11.2  |
| Males                |      |                         |       |
| 0.10                 | 1.0  | 0.8                     | 1.2   |
| 0.66                 | 2.0  | 1.9                     | 2.1   |
| 0.97                 | 3.0  | 2.8                     | 3.2   |
| 1.00                 | 4.0  | 3.6                     | 4.4   |
| 1.00                 | 5.0  | 4.5                     | 5.5   |
| 1.00                 | 6.0  | 5.3                     | 6.7   |
| 1.00                 | 7.0  | 6.1                     | 7.9   |
| 1.00                 | 8.0  | 7.0                     | 9.0   |
| 1.00                 | 9.0  | 7.8                     | 10.2  |
| 1.00                 | 10.0 | 8.7                     | 11.3  |
| 1.00                 | 11.0 | 9.5                     | 12.5  |
| 1.00                 | 12.0 | 10.3                    | 13.7  |

<sup>a</sup> Inshore strata 1–46, 52, 55, offshore strata 1–19, 61–76.

## Silver Hake: *Merluccius bilinearis*

Silver hake are widely distributed in the northwest Atlantic from Newfoundland to South Carolina, but are most abundant in coastal waters from Maine to New Jersey; two stocks inhabit this region, with Georges Bank as the boundary (Almeida 1987). Spawning of both stocks takes place from spring through early autumn with most activity occurring between May and September (Table 2). Total landings from both stocks combined declined from over 100,000 metric tons annually during the early 1970's, when distant water fleets were dominant, to less than 20,000 t since 1986 (NEFSC 1991).

Median length at maturity for female and male silver hake from the Gulf of Maine-northern Georges Bank stock was 23.1 and 22.3 cm, respectively (Tables 3 and 21, Fig. 19). For silver hake from the southern Georges Bank stock, values of  $L_{50}$  were 23.2 cm for females and 22.7 cm for males (Tables 3 and 23, Fig. 21). Median age at maturity for both sexes was 1.7 yr from the former region, and 1.6 yr from the latter region (Table

4, Figs. 20 and 22); proportions of mature fish calculated for each age are presented in Tables 22 and 24.

**Table 21**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Gulf of Maine-northern Georges Bank female ( $n=1410$ ) and male ( $n=1026$ ) silver hake, *Merluccius bilinearis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985-1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 17.4           | 16.6                    | 18.2  |
| 0.10                 | 18.8           | 18.2                    | 19.5  |
| 0.15                 | 19.7           | 19.2                    | 20.3  |
| 0.20                 | 20.4           | 19.9                    | 20.9  |
| 0.25                 | 20.9           | 20.5                    | 21.4  |
| 0.30                 | 21.4           | 21.0                    | 21.9  |
| 0.35                 | 21.9           | 21.5                    | 22.3  |
| 0.40                 | 22.3           | 21.9                    | 22.7  |
| 0.45                 | 22.7           | 22.3                    | 23.1  |
| 0.50                 | 23.1           | 22.7                    | 23.5  |
| 0.55                 | 23.5           | 23.1                    | 23.8  |
| 0.60                 | 23.8           | 23.5                    | 24.2  |
| 0.65                 | 24.3           | 23.9                    | 24.7  |
| 0.70                 | 24.7           | 24.3                    | 25.1  |
| 0.75                 | 25.2           | 24.8                    | 25.6  |
| 0.80                 | 25.7           | 25.3                    | 26.2  |
| 0.85                 | 26.4           | 25.9                    | 26.9  |
| 0.90                 | 27.3           | 26.7                    | 27.9  |
| 0.95                 | 28.7           | 28.0                    | 29.4  |
| 0.99                 | 31.9           | 30.9                    | 32.9  |
| Males                |                |                         |       |
| 0.05                 | 17.7           | 17.0                    | 18.4  |
| 0.10                 | 18.9           | 18.3                    | 19.4  |
| 0.15                 | 19.6           | 19.0                    | 20.1  |
| 0.20                 | 20.1           | 19.6                    | 20.6  |
| 0.25                 | 20.6           | 20.1                    | 21.0  |
| 0.30                 | 20.9           | 20.5                    | 21.4  |
| 0.35                 | 21.3           | 20.9                    | 21.7  |
| 0.40                 | 21.6           | 21.2                    | 22.0  |
| 0.45                 | 21.9           | 21.6                    | 22.3  |
| 0.50                 | 22.3           | 21.9                    | 22.6  |
| 0.55                 | 22.6           | 22.2                    | 22.9  |
| 0.60                 | 22.9           | 22.5                    | 23.3  |
| 0.65                 | 23.2           | 22.8                    | 23.6  |
| 0.70                 | 23.6           | 23.2                    | 24.0  |
| 0.75                 | 24.0           | 23.5                    | 24.4  |
| 0.80                 | 24.4           | 23.9                    | 24.9  |
| 0.85                 | 24.9           | 24.4                    | 25.4  |
| 0.90                 | 25.7           | 25.1                    | 26.2  |
| 0.95                 | 26.8           | 26.1                    | 27.5  |
| 0.99                 | 29.4           | 28.4                    | 30.3  |

<sup>a</sup> Offshore strata 20-30, 36-40.




**Figure 19**

Calculated and observed (•) proportion mature at length based on macroscopic observations of whole gonads of Gulf of Maine-northern Georges Bank female ( $n=1410$ ) and male ( $n=1026$ ) silver hake, *Merluccius bilinearis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985-1989.

Values of  $L_{50}$  above are 2–4 cm lower for both sexes than those reported by Morse (1979)<sup>2</sup> for the year 1977 over the entire region. Doubleday and Halliday (1975) observed that the median length at maturity for silver hake from the Scotian Shelf occurred at 26–27 cm in females and at 23–24 cm in males, with  $A_{50}$  values lying between

ages 1 and 2 for both sexes. Beacham (1983c) further noted a decline in  $L_{50}$  for females from 31 cm during the 1960's to 26 cm during the late 1970's in the same area. A similar decline from 27 cm to 24 cm was found for males during the same period, although  $A_{50}$  values generally remained between 1 and 2 yr for both sexes.




**Figure 20**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of Gulf of Maine-northern Georges Bank female ( $n=1410$ ) and male ( $n=1026$ ) silver hake, *Merluccius bilinearis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1989.

**Table 22**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Gulf of Maine-northern Georges Bank female ( $n=1410$ ) and male ( $n=1026$ ) silver hake *Merluccius bilinearis* sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Age | 95% Confidence interval |       |
|----------------------|-----|-------------------------|-------|
|                      |     | Lower                   | Upper |
| Females              |     |                         |       |
| 0.08                 | 1.0 | 0.9                     | 1.1   |
| 0.70                 | 2.0 | 1.9                     | 2.1   |
| 0.99                 | 3.0 | 2.9                     | 3.1   |
| 1.00                 | 4.0 | 3.8                     | 4.2   |
| 1.00                 | 5.0 | 4.7                     | 5.3   |
| 1.00                 | 6.0 | 5.6                     | 6.4   |
| 1.00                 | 7.0 | 6.5                     | 7.5   |
| 1.00                 | 8.0 | 7.4                     | 8.6   |
| 1.00                 | 9.0 | 8.3                     | 9.7   |
| Males                |     |                         |       |
| 0.07                 | 1.0 | 0.9                     | 1.1   |
| 0.74                 | 2.0 | 1.9                     | 2.1   |
| 0.99                 | 3.0 | 2.8                     | 3.2   |
| 1.00                 | 4.0 | 3.7                     | 4.3   |
| 1.00                 | 5.0 | 4.6                     | 5.4   |
| 1.00                 | 6.0 | 5.5                     | 6.5   |
| 1.00                 | 7.0 | 6.4                     | 7.6   |
| 1.00                 | 8.0 | 7.3                     | 8.7   |

<sup>a</sup> Offshore strata 20–30, 36–40.




**Figure 21**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of southern Georges Bank - Middle Atlantic female ( $n=2429$ ) and male ( $n=1612$ ) silver hake, *Merluccius bilinearis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1989.

**Table 23**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of southern Georges Bank - Middle Atlantic female ( $n=2429$ ) and male ( $n=1612$ ) silver hake, *Merluccius bilinearis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 16.8           | 16.0                    | 17.6  |
| 0.10                 | 18.4           | 17.7                    | 19.1  |
| 0.15                 | 19.4           | 18.8                    | 20.1  |
| 0.20                 | 20.2           | 19.6                    | 20.8  |
| 0.25                 | 20.8           | 20.3                    | 21.3  |
| 0.30                 | 21.4           | 20.9                    | 21.8  |
| 0.35                 | 21.8           | 21.4                    | 22.3  |
| 0.40                 | 22.3           | 21.9                    | 22.8  |
| 0.45                 | 22.7           | 22.3                    | 23.2  |
| 0.50                 | 23.2           | 22.8                    | 23.6  |
| 0.55                 | 23.6           | 23.2                    | 24.0  |
| 0.60                 | 24.1           | 23.7                    | 24.4  |
| 0.65                 | 24.5           | 24.1                    | 24.9  |
| 0.70                 | 25.0           | 24.6                    | 25.4  |
| 0.75                 | 25.5           | 25.2                    | 25.9  |
| 0.80                 | 26.2           | 25.8                    | 26.6  |
| 0.85                 | 26.9           | 26.5                    | 27.3  |
| 0.90                 | 27.9           | 27.4                    | 28.4  |
| 0.95                 | 29.5           | 29.0                    | 30.1  |
| 0.99                 | 33.1           | 32.2                    | 33.9  |
| Males                |                |                         |       |
| 0.05                 | 18.0           | 17.2                    | 18.8  |
| 0.10                 | 19.2           | 18.5                    | 19.9  |
| 0.15                 | 19.9           | 19.3                    | 20.5  |
| 0.20                 | 20.5           | 19.9                    | 21.0  |
| 0.25                 | 20.9           | 20.4                    | 21.4  |
| 0.30                 | 21.3           | 20.9                    | 21.8  |
| 0.35                 | 21.7           | 21.3                    | 22.1  |
| 0.40                 | 22.0           | 21.6                    | 22.5  |
| 0.45                 | 22.4           | 22.0                    | 22.8  |
| 0.50                 | 22.7           | 22.3                    | 23.1  |
| 0.55                 | 23.0           | 22.6                    | 23.4  |
| 0.60                 | 23.3           | 23.0                    | 23.7  |
| 0.65                 | 23.7           | 23.3                    | 24.0  |
| 0.70                 | 24.0           | 23.7                    | 24.4  |
| 0.75                 | 24.4           | 24.1                    | 24.8  |
| 0.80                 | 24.9           | 24.5                    | 25.2  |
| 0.85                 | 25.4           | 25.1                    | 25.8  |
| 0.90                 | 26.2           | 25.8                    | 26.6  |
| 0.95                 | 27.4           | 26.9                    | 27.9  |
| 0.99                 | 30.0           | 29.2                    | 30.7  |

<sup>a</sup> Inshore strata 1–46, 52, 55, offshore strata 1–19, 61–76.




**Figure 22**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of southern Georges Bank - Middle Atlantic female ( $n=2429$ ) and male ( $n=1612$ ) silver hake, *Merluccius bilinearis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1989.

**Table 24**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of southern Georges Bank - Middle Atlantic female ( $n=2429$ ) and male ( $n=1612$ ) silver hake, *Merluccius bilinearis*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Age | 95% Confidence interval |       |
|----------------------|-----|-------------------------|-------|
|                      |     | Lower                   | Upper |
| Females              |     |                         |       |
| 0.10                 | 1.0 | 0.9                     | 1.1   |
| 0.81                 | 2.0 | 2.0                     | 2.0   |
| 0.99                 | 3.0 | 2.9                     | 3.1   |
| 1.00                 | 4.0 | 3.8                     | 4.2   |
| 1.00                 | 5.0 | 4.7                     | 5.3   |
| 1.00                 | 6.0 | 5.6                     | 6.4   |
| 1.00                 | 7.0 | 6.6                     | 7.4   |
| Males                |     |                         |       |
| 0.08                 | 1.0 | 0.9                     | 1.1   |
| 0.79                 | 2.0 | 1.9                     | 2.1   |
| 0.99                 | 3.0 | 2.9                     | 3.1   |
| 1.00                 | 4.0 | 3.8                     | 4.2   |
| 1.00                 | 5.0 | 4.7                     | 5.3   |
| 1.00                 | 6.0 | 5.6                     | 6.4   |
| 1.00                 | 7.0 | 6.5                     | 7.5   |
| 1.00                 | 8.0 | 7.4                     | 8.6   |
| 1.00                 | 9.0 | 8.3                     | 9.7   |

<sup>a</sup> Inshore strata 1–46, 52, 55, offshore strata 1–19, 61–76.

## Redfish: *Sebastes fasciatus*

Redfish occur throughout the north Atlantic from Norway to Georges Bank; off New England, *S. fasciatus* are most abundant in deeper waters of the Gulf of Maine where a single stock is thought to exist (Mayo 1980). Reproduction is viviparous and mating takes place in autumn when most males are mature; larval extrusion occurs during the following spring and summer (Table 2). Total commercial landings of this species have declined from a peak of 60,000 metric tons per year in the early 1940's to about 15,000 t in the late 1970's; total landings have declined further to less than 2,000 t per year since 1987 (NEFSC 1991).

Median length at maturity for female and male redfish from the Gulf of Maine was 22.3 and 20.9 cm, respectively (Tables 3 and 25, Fig. 23). Median age at maturity was 5.5 yr for both sexes (Table 4, Fig. 24); proportions of mature fish calculated for each age are presented in Table 26. These results, extracted from Mayo et al. (1990), are representative of the years 1975–

1980, the most recent period for which a consistent series of maturation data are available from the Northeast Fisheries Science Center survey. Because of the




**Figure 23**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=1457$ ) and male ( $n=1526$ ) redfish, *Sebastes fasciatus*, from the Gulf of Maine sampled, respectively, during Northeast Fisheries Science Center spring and autumn bottom trawl surveys, 1975–1980.

**Table 25**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=1457$ ) and male ( $n=1526$ ) redfish, *Sebastes fasciatus*, from the Gulf of Maine sampled, respectively, during Northeast Fisheries Science Center spring and autumn bottom trawl surveys<sup>a</sup>, 1975–1980.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 17.3           | 16.4                    | 18.2  |
| 0.10                 | 18.6           | 17.8                    | 19.3  |
| 0.15                 | 19.3           | 18.7                    | 20.0  |
| 0.20                 | 19.9           | 19.3                    | 20.5  |
| 0.25                 | 20.4           | 19.9                    | 20.9  |
| 0.30                 | 20.8           | 20.3                    | 21.3  |
| 0.35                 | 21.2           | 20.8                    | 21.7  |
| 0.40                 | 21.6           | 21.1                    | 22.0  |
| 0.45                 | 21.9           | 21.5                    | 22.3  |
| 0.50                 | 22.3           | 21.9                    | 22.7  |
| 0.55                 | 22.6           | 22.2                    | 23.0  |
| 0.60                 | 22.9           | 22.5                    | 23.3  |
| 0.65                 | 23.3           | 22.9                    | 23.7  |
| 0.70                 | 23.7           | 23.3                    | 24.1  |
| 0.75                 | 24.1           | 23.7                    | 24.5  |
| 0.80                 | 24.6           | 24.1                    | 25.0  |
| 0.85                 | 25.2           | 24.6                    | 25.7  |
| 0.90                 | 25.9           | 25.3                    | 26.5  |
| 0.95                 | 27.2           | 26.5                    | 27.9  |
| 0.99                 | 30.0           | 28.9                    | 31.1  |
| Males                |                |                         |       |
| 0.05                 | 12.0           | 10.6                    | 13.5  |
| 0.10                 | 14.3           | 13.1                    | 15.5  |
| 0.15                 | 15.7           | 14.6                    | 16.7  |
| 0.20                 | 16.7           | 15.8                    | 17.7  |
| 0.25                 | 17.6           | 16.8                    | 18.5  |
| 0.30                 | 18.4           | 17.6                    | 19.2  |
| 0.35                 | 19.1           | 18.3                    | 19.8  |
| 0.40                 | 19.7           | 19.1                    | 20.4  |
| 0.45                 | 20.3           | 19.7                    | 20.9  |
| 0.50                 | 20.9           | 20.4                    | 21.5  |
| 0.55                 | 21.5           | 21.0                    | 22.1  |
| 0.60                 | 22.2           | 21.7                    | 22.7  |
| 0.65                 | 22.8           | 22.3                    | 23.3  |
| 0.70                 | 23.5           | 23.0                    | 24.0  |
| 0.75                 | 24.3           | 23.8                    | 24.7  |
| 0.80                 | 25.1           | 24.6                    | 25.6  |
| 0.85                 | 26.2           | 25.6                    | 26.7  |
| 0.90                 | 27.6           | 26.9                    | 28.2  |
| 0.95                 | 29.8           | 29.0                    | 30.7  |
| 0.99                 | 34.8           | 33.4                    | 36.2  |

<sup>a</sup> Offshore strata 24, 26–30, 36–40.

disparity in seasonal maturation between sexes related to redfish's viviparous reproduction, analyses were performed on spring data for females and autumn data for males.

Values of  $L_{50}$  above are comparable to those for both sexes reported by Morse (1979)<sup>2</sup> for the year 1977. However considerable variability in size and age at maturation in relation to early growth and environmental conditions was noted by Mayo et al. (1990). Substantial differences between sexes were observed on the southern Scotian Shelf where Ni and Sandeman (1984) reported  $L_{50}$  values of 25.9 and 15.9 cm for *S. fasciatus* females and males, respectively. Information of growth of immature redfish presented by Perlmutter and Clarke (1949) suggested that the median age at maturation in the 1940's was likely to have been in the range of 8–9 yr, in contrast to the more recent value of 5.5 yr noted above.




**Figure 24**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=1457$ ) and male ( $n=1526$ ) redfish, *Sebastes fasciatus*, from the Gulf of Maine sampled during Northeast Fisheries Science Center spring and autumn bottom trawl surveys, 1975–1980.

**Table 26**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=1457$ ) and male ( $n=1526$ ) redfish, *Sebastes fasciatus*, from the Gulf of Maine sampled during Northeast Fisheries Science Center spring and autumn bottom trawl surveys<sup>a</sup>, 1975–1980.

| Proportion<br>mature | Age               | 95% Confidence interval |       |
|----------------------|-------------------|-------------------------|-------|
|                      |                   | Lower                   | Upper |
| Females              |                   |                         |       |
| Spring               |                   |                         |       |
| 0.01                 | 1.0               | 0.2                     | 1.8   |
| 0.02                 | 2.0               | 1.4                     | 2.6   |
| 0.05                 | 3.0               | 2.5                     | 3.5   |
| 0.15                 | 4.0               | 3.7                     | 4.3   |
| 0.36                 | 5.0               | 4.8                     | 5.2   |
| 0.64                 | 6.0               | 5.8                     | 6.2   |
| 0.85                 | 7.0               | 6.7                     | 7.3   |
| 0.95                 | 8.0               | 7.6                     | 8.4   |
| 0.98                 | 9.0               | 8.5                     | 9.5   |
| 0.99                 | 10.0              | 9.3                     | 10.7  |
| 1.00                 | 11.0              | 10.1                    | 11.9  |
| 1.00                 | 12.0              | 11.0                    | 13.0  |
| 1.00                 | 13.0              | 11.8                    | 14.2  |
| 1.00                 | 14.0              | 12.7                    | 15.3  |
| 1.00                 | 15.0              | 13.5                    | 16.5  |
| 1.00                 | 16.0              | 14.4                    | 17.6  |
| 1.00                 | 17.0              | 15.2                    | 18.8  |
| 1.00                 | 18.0              | 16.0                    | 20.0  |
| 1.00                 | 19.0              | 16.9                    | 21.1  |
| 1.00                 | 20.0 <sup>b</sup> | 17.7                    | 22.3  |
| Males                |                   |                         |       |
| Autumn               |                   |                         |       |
| 0.06                 | 1.0               | 0.1                     | 1.9   |
| 0.11                 | 2.0               | 1.3                     | 2.7   |
| 0.18                 | 3.0               | 2.4                     | 3.6   |
| 0.28                 | 4.0               | 3.6                     | 4.4   |
| 0.42                 | 5.0               | 4.7                     | 5.3   |
| 0.57                 | 6.0               | 5.7                     | 6.3   |
| 0.71                 | 7.0               | 6.7                     | 7.3   |
| 0.81                 | 8.0               | 7.6                     | 8.4   |
| 0.89                 | 9.0               | 8.5                     | 9.5   |
| 0.94                 | 10.0              | 9.3                     | 10.7  |
| 0.96                 | 11.0              | 10.2                    | 11.8  |
| 0.98                 | 12.0              | 11.0                    | 13.0  |
| 0.99                 | 13.0              | 11.9                    | 14.1  |
| 0.99                 | 14.0              | 12.7                    | 15.3  |
| 1.00                 | 15.0              | 13.6                    | 16.4  |
| 1.00                 | 16.0              | 14.4                    | 17.6  |
| 1.00                 | 17.0              | 15.2                    | 18.8  |
| 1.00                 | 18.0              | 16.1                    | 19.9  |
| 1.00                 | 19.0              | 16.9                    | 21.1  |
| 1.00                 | 20.0 <sup>b</sup> | 17.7                    | 22.3  |

<sup>a</sup> Offshore strata 24, 26–30, 36–40.

<sup>b</sup> Older ages (21–53) are 100% mature.

Scup: *Stenotomus chrysops*

Scup range from the Gulf of Maine to Cape Hatteras, North Carolina (Bigelow and Schroeder 1953). Spawning occurs in coastal waters from May to August with peak spawning during May and June (Table 2). Commercial landings of scup were between 4,000 and 5,000 metric tons in the early 1970's and by 1980 had increased to about 12,000 t, subsequently declining to about 7,000 t in 1989 (NEFSC 1991).

Median length at maturity was 15.5 cm and 15.6 cm for females and males, respectively, collected during the spring of 1985, and 1987–1990 (Tables 3 and 27, Fig. 25).

Similar values were calculated for scup in the Peconic Bays of eastern Long Island during 1964–1966. Both female and male scup were determined to be sexually mature at age two (Finkelstein 1969b), which corresponded to a length of about 16 cm for both sexes (Finkelstein 1969a).




Figure 25

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=284$ ) and male ( $n=232$ ) scup, *Stenotomus chrysops*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985, 1987–1990.

Table 27

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=284$ ) and male ( $n=232$ ) scup, *Stenotomus chrysops*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985, 1987–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 12.3           | 11.2                    | 13.4  |
| 0.10                 | 13.1           | 12.2                    | 14.0  |
| 0.15                 | 13.6           | 12.8                    | 14.4  |
| 0.20                 | 14.0           | 13.2                    | 14.7  |
| 0.25                 | 14.3           | 13.6                    | 15.0  |
| 0.30                 | 14.6           | 13.9                    | 15.2  |
| 0.35                 | 14.8           | 14.2                    | 15.5  |
| 0.40                 | 15.1           | 14.5                    | 15.7  |
| 0.45                 | 15.3           | 14.7                    | 15.9  |
| 0.50                 | 15.5           | 14.9                    | 16.1  |
| 0.55                 | 15.7           | 15.2                    | 16.3  |
| 0.60                 | 16.0           | 15.4                    | 16.5  |
| 0.65                 | 16.2           | 15.6                    | 16.8  |
| 0.70                 | 16.4           | 15.8                    | 17.0  |
| 0.75                 | 16.7           | 16.1                    | 17.4  |
| 0.80                 | 17.0           | 16.4                    | 17.7  |
| 0.85                 | 17.4           | 16.7                    | 18.1  |
| 0.90                 | 17.9           | 17.1                    | 18.7  |
| 0.95                 | 18.8           | 17.8                    | 19.7  |
| 0.99                 | 20.6           | 19.2                    | 22.0  |
| Males                |                |                         |       |
| 0.05                 | 11.9           | 10.6                    | 13.2  |
| 0.10                 | 12.9           | 11.8                    | 13.9  |
| 0.15                 | 13.4           | 12.5                    | 14.4  |
| 0.20                 | 13.9           | 13.0                    | 14.8  |
| 0.25                 | 14.2           | 13.4                    | 15.1  |
| 0.30                 | 14.5           | 13.8                    | 15.3  |
| 0.35                 | 14.8           | 14.1                    | 15.6  |
| 0.40                 | 15.1           | 14.4                    | 15.8  |
| 0.45                 | 15.4           | 14.6                    | 16.1  |
| 0.50                 | 15.6           | 14.9                    | 16.3  |
| 0.55                 | 15.9           | 15.1                    | 16.6  |
| 0.60                 | 16.1           | 15.4                    | 16.8  |
| 0.65                 | 16.4           | 15.7                    | 17.1  |
| 0.70                 | 16.7           | 15.9                    | 17.4  |
| 0.75                 | 17.0           | 16.2                    | 17.8  |
| 0.80                 | 17.3           | 16.5                    | 18.2  |
| 0.85                 | 17.8           | 16.9                    | 18.7  |
| 0.90                 | 18.3           | 17.4                    | 19.3  |
| 0.95                 | 19.3           | 18.1                    | 20.5  |
| 0.99                 | 21.3           | 19.7                    | 23.0  |

<sup>a</sup> Offshore strata 2, 3, 61–63, 65–67, 70–71, 74–75.

## Black Sea Bass: *Centropristis striata*

Black sea bass range from Cape Cod, Massachusetts, to northern Florida, but are most abundant in the Mid-Atlantic region. Meristic and morphometric studies indicate that only one stock exists north of Cape Hatteras, North Carolina (Shepherd 1991). Spawning of this stock progresses northward from March off North Carolina to October in Cape Cod waters (NEFSC 1991). Commercial landings of black sea bass have averaged 1,500 metric tons per year during 1980–1989 while annual recreational landings have fluctuated between 500 t and 8,100 t over the same period (NEFSC 1991).

Median length at maturity for female and male black sea bass was 19.1 and 19.0 cm (TL), respectively (Tables 3 and 28, Fig. 26). These sizes occur between ages 2 and 3 according to growth functions developed by Mercer (1978) for black sea bass north and south of Cape Hatteras and by Wenner et al. (1986) for black sea bass in the South Atlantic. Cupka et al. (1973)<sup>4</sup> reported that both sexes matured between 14 and 18 cm (standard length) in the South Carolina region. Wenner et al. (1986) reported that most females mature at age 2; Mercer (1978) observed mature males in all age groups, including age 1. Maturity analyses are complicated by

the fact that black sea bass are protogynous hermaphrodites (Lavenda 1949), and that the proportion of mature males obtained through sex succession of females is still uncertain (Mercer 1978; Wenner et al. 1986).

<sup>4</sup> D.M. Cupka, R.K. Dias and J. Tucker. 1973. Biology of the black sea bass, *Centropristis striata*, from South Carolina waters. Unpubl. manuscript. South Carolina Wildlife and Marine Resources Department, P.O. Box 12559, Charleston, SC 29412, 93 p.

**Table 28**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=561$ ) and male ( $n=348$ ) black sea bass, *Centropristis striata*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 7.1            | 4.2                     | 10.1  |
| 0.10                 | 10.2           | 7.8                     | 12.6  |
| 0.15                 | 12.1           | 10.0                    | 14.2  |
| 0.20                 | 13.5           | 11.6                    | 15.3  |
| 0.25                 | 14.7           | 13.0                    | 16.3  |
| 0.30                 | 15.7           | 14.2                    | 17.2  |
| 0.35                 | 16.6           | 15.3                    | 18.0  |
| 0.40                 | 17.5           | 16.3                    | 18.7  |
| 0.45                 | 18.3           | 17.2                    | 19.5  |
| 0.50                 | 19.1           | 18.1                    | 20.2  |
| 0.55                 | 20.0           | 19.0                    | 20.9  |
| 0.60                 | 20.8           | 19.9                    | 21.7  |
| 0.65                 | 21.7           | 20.8                    | 22.6  |
| 0.70                 | 22.6           | 21.7                    | 23.5  |
| 0.75                 | 23.6           | 22.7                    | 24.6  |
| 0.80                 | 24.8           | 23.7                    | 25.9  |
| 0.85                 | 26.2           | 25.0                    | 27.5  |
| 0.90                 | 28.1           | 26.6                    | 29.6  |
| 0.95                 | 31.2           | 29.2                    | 33.2  |
| 0.99                 | 37.9           | 34.7                    | 41.1  |
| Males                |                |                         |       |
| 0.05                 | 10.8           | 8.2                     | 13.5  |
| 0.10                 | 12.9           | 10.7                    | 15.1  |
| 0.15                 | 14.2           | 12.3                    | 16.1  |
| 0.20                 | 15.1           | 13.4                    | 16.8  |
| 0.25                 | 15.9           | 14.4                    | 17.5  |
| 0.30                 | 16.6           | 15.2                    | 18.1  |
| 0.35                 | 17.3           | 16.0                    | 18.6  |
| 0.40                 | 17.9           | 16.6                    | 19.1  |
| 0.45                 | 18.4           | 17.3                    | 19.6  |
| 0.50                 | 19.0           | 17.9                    | 20.1  |
| 0.55                 | 19.5           | 18.5                    | 20.6  |
| 0.60                 | 20.1           | 19.1                    | 21.1  |
| 0.65                 | 20.7           | 19.7                    | 21.7  |
| 0.70                 | 21.3           | 20.4                    | 22.3  |
| 0.75                 | 22.0           | 21.0                    | 23.1  |
| 0.80                 | 22.8           | 21.7                    | 23.9  |
| 0.85                 | 23.8           | 22.6                    | 25.0  |
| 0.90                 | 25.1           | 23.7                    | 26.5  |
| 0.95                 | 27.2           | 25.3                    | 29.0  |
| 0.99                 | 31.7           | 28.9                    | 34.6  |

<sup>a</sup> Inshore strata 1–61, offshore strata 1–12, 25, 61–76.




**Figure 26**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=561$ ) and male ( $n=348$ ) black sea bass, *Centropristis striata*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

## Ocean Pout: *Macrozoarces americanus*

Ocean pout are distributed from the Gulf of St. Lawrence to Delaware (Olsen and Merriman 1946); off the coast of New England they are abundant in the Gulf of Maine and Southern New England area. Spawning occurs in rocky areas from September to October (Table 2). Landings of ocean pout fluctuated about an average of 600 metric tons during 1975–1983, then increased to a peak of about 2,200 t in 1987 and have recently declined to about 1,300 t (NEFSC 1991).

Median length at maturity for female and male ocean pout from the Gulf of Maine area was 26.2 cm and 30.3 cm, respectively (Tables 3 and 29, Fig. 27). Ocean pout from the Southern New England area had  $L_{50}$  values of 31.3 cm and 31.9 cm for females and males, respectively (Tables 3 and 30, Fig. 28).

Larger  $L_{50}$  values were calculated by Olsen and Merriman (1946) for ocean pout taken in the Southern New England area in May of 1944. Females were found to be mature at an average length of 55 cm which

corresponded to an age of either 6 or 7 yr. All male ocean pout below 25 cm were immature and nearly all




**Figure 27**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=88$ ) and male ( $n=55$ ) ocean pout, *Macrozoarces americanus*, from the Gulf of Maine sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 29**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=88$ ) and male ( $n=55$ ) ocean pout, *Macrozoarces americanus*, from the Gulf of Maine sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 11.1           | -3.3                    | 25.5  |
| 0.10                 | 14.9           | 2.9                     | 26.9  |
| 0.15                 | 17.3           | 6.8                     | 27.8  |
| 0.20                 | 19.1           | 9.6                     | 28.5  |
| 0.25                 | 20.6           | 12.0                    | 29.2  |
| 0.30                 | 21.8           | 14.0                    | 29.7  |
| 0.35                 | 23.0           | 15.8                    | 30.3  |
| 0.40                 | 24.1           | 17.4                    | 30.8  |
| 0.45                 | 25.2           | 19.0                    | 31.3  |
| 0.50                 | 26.2           | 20.5                    | 31.9  |
| 0.55                 | 27.2           | 22.0                    | 32.5  |
| 0.60                 | 28.3           | 23.4                    | 33.2  |
| 0.65                 | 29.4           | 24.8                    | 34.0  |
| 0.70                 | 30.5           | 26.2                    | 34.9  |
| 0.75                 | 31.8           | 27.5                    | 36.1  |
| 0.80                 | 33.3           | 28.9                    | 37.7  |
| 0.85                 | 35.1           | 30.3                    | 39.9  |
| 0.90                 | 37.5           | 31.8                    | 43.2  |
| 0.95                 | 41.3           | 33.7                    | 48.9  |
| 0.99                 | 49.8           | 37.1                    | 62.5  |
| Males                |                |                         |       |
| 0.05                 | 10.3           | -13.8                   | 34.4  |
| 0.10                 | 15.4           | -4.9                    | 35.7  |
| 0.15                 | 18.5           | 0.5                     | 36.5  |
| 0.20                 | 20.9           | 4.6                     | 37.1  |
| 0.25                 | 22.8           | 8.0                     | 37.7  |
| 0.30                 | 24.6           | 10.9                    | 38.2  |
| 0.35                 | 26.1           | 13.5                    | 38.7  |
| 0.40                 | 27.6           | 16.0                    | 39.2  |
| 0.45                 | 29.0           | 18.3                    | 39.7  |
| 0.50                 | 30.3           | 20.5                    | 40.2  |
| 0.55                 | 31.7           | 22.6                    | 40.8  |
| 0.60                 | 33.1           | 24.8                    | 41.4  |
| 0.65                 | 34.5           | 26.9                    | 42.2  |
| 0.70                 | 36.1           | 29.1                    | 43.1  |
| 0.75                 | 37.8           | 31.3                    | 44.4  |
| 0.80                 | 39.8           | 33.4                    | 46.1  |
| 0.85                 | 42.1           | 35.6                    | 48.7  |
| 0.90                 | 45.3           | 37.7                    | 52.9  |
| 0.95                 | 50.4           | 40.1                    | 60.7  |
| 0.99                 | 61.6           | 43.5                    | 79.7  |

<sup>a</sup> Inshore strata 57–66, offshore strata 26–40.

males greater than 39 cm were mature at an age of 3 or 4 yr. A study conducted by Keats et al. (1985) in Newfoundland waters during July and August, 1983–1984, reported that the mean lengths of nesting pairs of ocean pout were 70.5 cm and 81.1 cm for females and males, respectively.

The difference in  $L_{50}$  values of females in the present study compared to that of Olsen and Merriman (1946) may be partly due to difficulty in identifying a mature female. Larger females appear to have three different sizes of eggs present while developing; however, Keats et al. (1985) reported that egg masses guarded by females were uniform and appeared to be from one spawning, and concluded that multiple spawning was unlikely. Possibly, immature females develop eggs each

year for three years, spawning in the third year. This maturation process would lead to misclassification of immature fish as developing for two years, therefore artificially lowering the  $L_{50}$  value.




**Figure 28**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=197$ ) and male ( $n=205$ ) ocean pout, *Macrozoarces americanus*, from Southern New England sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 30**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=197$ ) and male ( $n=205$ ) ocean pout, *Macrozoarces americanus*, from Southern New England sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 17.1           | 8.4                     | 25.8  |
| 0.10                 | 20.7           | 13.3                    | 28.2  |
| 0.15                 | 22.9           | 16.3                    | 29.6  |
| 0.20                 | 24.6           | 18.5                    | 30.7  |
| 0.25                 | 26.0           | 20.3                    | 31.7  |
| 0.30                 | 27.2           | 21.9                    | 32.5  |
| 0.35                 | 28.3           | 23.3                    | 33.3  |
| 0.40                 | 29.3           | 24.6                    | 34.0  |
| 0.45                 | 30.3           | 25.9                    | 34.7  |
| 0.50                 | 31.3           | 27.1                    | 35.5  |
| 0.55                 | 32.2           | 28.2                    | 36.2  |
| 0.60                 | 33.2           | 29.4                    | 37.0  |
| 0.65                 | 34.2           | 30.6                    | 37.8  |
| 0.70                 | 35.3           | 31.9                    | 38.8  |
| 0.75                 | 36.5           | 33.2                    | 39.9  |
| 0.80                 | 37.9           | 34.6                    | 41.2  |
| 0.85                 | 39.6           | 36.3                    | 42.9  |
| 0.90                 | 41.8           | 38.3                    | 45.3  |
| 0.95                 | 45.4           | 41.1                    | 49.7  |
| 0.99                 | 53.3           | 46.7                    | 60.0  |
| Males                |                |                         |       |
| 0.05                 | 16.0           | 8.4                     | 23.7  |
| 0.10                 | 20.0           | 13.6                    | 26.5  |
| 0.15                 | 22.5           | 16.8                    | 28.3  |
| 0.20                 | 24.4           | 19.2                    | 29.6  |
| 0.25                 | 26.0           | 21.2                    | 30.7  |
| 0.30                 | 27.3           | 22.9                    | 31.8  |
| 0.35                 | 28.5           | 24.4                    | 32.7  |
| 0.40                 | 29.7           | 25.8                    | 33.6  |
| 0.45                 | 30.8           | 27.1                    | 34.5  |
| 0.50                 | 31.9           | 28.4                    | 35.4  |
| 0.55                 | 32.9           | 29.6                    | 36.3  |
| 0.60                 | 34.1           | 30.9                    | 37.2  |
| 0.65                 | 35.2           | 32.1                    | 38.3  |
| 0.70                 | 36.4           | 33.4                    | 39.5  |
| 0.75                 | 37.8           | 34.8                    | 40.8  |
| 0.80                 | 39.3           | 36.2                    | 42.4  |
| 0.85                 | 41.2           | 37.9                    | 44.5  |
| 0.90                 | 43.7           | 40.0                    | 47.4  |
| 0.95                 | 47.7           | 43.1                    | 52.4  |
| 0.99                 | 56.6           | 49.4                    | 63.8  |

<sup>a</sup> Inshore strata 1–52, offshore strata 1–12, 61–76.

## Yellowtail Flounder: *Pleuronectes ferrugineus*

Yellowtail flounder range from Labrador to Virginia with three major stocks occurring off the coast of New England: Georges Bank, Cape Cod, and Southern New England (Royce et al. 1959; Lux 1963). The spawning season is from April to August with peak spawning occurring in May and June (Table 2). Commercial landings for the Georges Bank stock declined from about 15,000 metric tons in the early 1970's to about 4,500 t by 1978, increased to over 11,000 t in 1983, then subsequently declined to a record low of 1,100 t in 1989 (NEFSC 1991). Landings in 1990 were about 12,000 t, attributable to a strong 1987 year class. Landings of the Cape Cod stock increased from about 1,500 t in the early 1970's to a peak of about 5,000 t by 1980, then subsequently declined to about 900 t by 1989 (NEFSC 1991). Commercial landings of the Southern New England stock declined from a high of about 32,000 t in 1969 to a low of 1,700 in 1976, increased to a peak of about 18,000 t in 1983 and subsequently declined to a low of

about 900 t in 1988. Landings increased in 1989 to 2,500 t owing to the recruiting 1987 year class (NEFSC 1991).




**Figure 29**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of Georges Bank female ( $n=167$ ) and male ( $n=271$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 31**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Georges Bank female ( $n=167$ ) and male ( $n=271$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 20.8           | 15.7                    | 25.9  |
| 0.10                 | 22.1           | 17.7                    | 26.4  |
| 0.15                 | 22.8           | 18.9                    | 26.8  |
| 0.20                 | 23.4           | 19.9                    | 27.0  |
| 0.25                 | 23.9           | 20.6                    | 27.3  |
| 0.30                 | 24.4           | 21.3                    | 27.5  |
| 0.35                 | 24.8           | 21.9                    | 27.6  |
| 0.40                 | 25.1           | 22.4                    | 27.8  |
| 0.45                 | 25.5           | 22.9                    | 28.0  |
| 0.50                 | 25.8           | 23.4                    | 28.2  |
| 0.55                 | 26.1           | 23.9                    | 28.4  |
| 0.60                 | 26.5           | 24.4                    | 28.6  |
| 0.65                 | 26.9           | 24.9                    | 28.8  |
| 0.70                 | 27.3           | 25.4                    | 29.1  |
| 0.75                 | 27.7           | 26.0                    | 29.4  |
| 0.80                 | 28.2           | 26.5                    | 29.8  |
| 0.85                 | 28.8           | 27.2                    | 30.4  |
| 0.90                 | 29.6           | 27.9                    | 31.2  |
| 0.95                 | 30.8           | 28.8                    | 32.9  |
| 0.99                 | 33.6           | 30.2                    | 37.0  |
| Males                |                |                         |       |
| 0.05                 | 13.7           | 6.1                     | 21.2  |
| 0.10                 | 15.6           | 9.0                     | 22.2  |
| 0.15                 | 16.8           | 10.8                    | 22.9  |
| 0.20                 | 17.7           | 12.1                    | 23.3  |
| 0.25                 | 18.5           | 13.3                    | 23.7  |
| 0.30                 | 19.2           | 14.2                    | 24.1  |
| 0.35                 | 19.8           | 15.1                    | 24.4  |
| 0.40                 | 20.3           | 15.9                    | 24.7  |
| 0.45                 | 20.9           | 16.7                    | 25.0  |
| 0.50                 | 21.4           | 17.5                    | 25.3  |
| 0.55                 | 21.9           | 18.2                    | 25.6  |
| 0.60                 | 22.4           | 19.0                    | 25.9  |
| 0.65                 | 23.0           | 19.8                    | 26.2  |
| 0.70                 | 23.6           | 20.6                    | 26.6  |
| 0.75                 | 24.3           | 21.5                    | 27.0  |
| 0.80                 | 25.0           | 22.5                    | 27.5  |
| 0.85                 | 25.9           | 23.7                    | 28.2  |
| 0.90                 | 27.1           | 25.1                    | 29.2  |
| 0.95                 | 29.1           | 27.1                    | 31.1  |
| 0.99                 | 33.4           | 30.1                    | 36.8  |

<sup>a</sup> Offshore strata 13–21.

Median length at maturity for female and male yellowtail flounder sampled during spring 1985–1990 was, respectively, 25.8 cm and 21.4 cm for the Georges Bank stock (Tables 3 and 31, Fig. 29), 27.3 cm and 26.8 cm for the Cape Cod stock (Tables 3 and 33, Fig. 31), and 25.5 cm and 19.6 cm for the Southern New England stock (Tables 3 and 35, Fig. 33). Median age at maturity for female and male Georges Bank yellowtail flounder was 1.8 yr and 1.3 yr, respectively (Table 4, Fig. 30). For the Cape Cod stock,  $A_{50}$  values were 2.6 yr for both sexes (Table 4, Fig. 32). The Southern New England stock had  $A_{50}$  values of 1.6 yr for females and 1.8 yr for males (Table 4, Fig. 34). Proportions of mature fish at age are presented in Tables 32, 34, and 36.

Morse and Morris (1981)<sup>5</sup> calculated similar  $L_{50}$  values of 27.3 cm and 24.2 cm for female and male yellowtail flounder collected east of 72° long. during 1975–1980. Yellowtail flounder collected over the entire re-

gion during 1977 had  $L_{50}$  values of 27.5 cm and 24.3 cm for females and males, respectively (Morse 1979)<sup>2</sup>. Scott (1954) examined yellowtail flounder collected during July–September 1946 between Nantucket Shoals and Montauk Point, and determined that there were no immature females over 32 cm, and that 100% maturation was reached by age 3. Royce et al. (1959) determined from Southern New England samples collected in May 1943 that  $L_{50}$  for females was 32 cm and that males became sexually mature before recruitment to the commercial catch, at about 26 cm. Female yellowtail had  $A_{50}$  values of about 2 yr and males probably matured at about age 1.

Median length at maturity of yellowtail flounder collected during July in the Scotian Shelf area was 24 cm during 1970–1974 for females and increased to 29 cm during 1975–1979; for males,  $L_{50}$  remained at about 19–20 cm during the same time periods (Beacham 1983d). Values of  $A_{50}$  declined for females and males from 3.1 yr and 3.2 yr, respectively, during 1970–1974 to 2.9 yr and 2.0 yr, respectively, during 1975–1979 (Beacham 1983d).

<sup>5</sup> W.W. Morse and A. Morris. 1981. An analysis of yellowtail flounder, *Limanda ferruginea*, length at maturity east and west of 72° long. for 1975–1980. NMFS/NEFC, Sandy Hook Lab. Rep. No. 81-16, 4 p.




**Figure 30**


Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of Georges Bank female ( $n=167$ ) and male ( $n=271$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 32**

Calculated proportion mature at age with approximate 95 % confidence intervals based on macroscopic observations of whole gonads of Georges Bank female ( $n=167$ ) and male ( $n=271$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Age | 95% Confidence interval |       |
|----------------------|-----|-------------------------|-------|
|                      |     | Lower                   | Upper |
| Females              |     |                         |       |
| 0.00                 | 1.0 | -2.1                    | 4.1   |
| 0.88                 | 2.0 | 1.9                     | 2.1   |
| 1.00                 | 3.0 | -0.1                    | 6.1   |
| 1.00                 | 4.0 | -2.1                    | 10.1  |
| 1.00                 | 5.0 | -4.2                    | 14.2  |
| 1.00                 | 6.0 | -6.3                    | 18.3  |
| 1.00                 | 7.0 | -8.3                    | 22.3  |
| Males                |     |                         |       |
| 0.22                 | 1.0 | 0.4                     | 1.6   |
| 0.93                 | 2.0 | 1.8                     | 2.2   |
| 1.00                 | 3.0 | 2.4                     | 3.6   |
| 1.00                 | 4.0 | 2.8                     | 5.2   |
| 1.00                 | 5.0 | 3.2                     | 6.8   |
| 1.00                 | 6.0 | 3.7                     | 8.3   |
| 1.00                 | 7.0 | 4.1                     | 9.9   |

<sup>a</sup> Offshore strata 13–21.

**Figure 31**


Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of Cape Cod female ( $n=140$ ) and male ( $n=165$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 33**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Cape Cod female ( $n=140$ ) and male ( $n=165$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Female               |                |                         |       |
| 0.05                 | 20.5           | 17.8                    | 23.1  |
| 0.10                 | 22.2           | 20.0                    | 24.4  |
| 0.15                 | 23.3           | 21.3                    | 25.2  |
| 0.20                 | 24.1           | 22.3                    | 25.9  |
| 0.25                 | 24.7           | 23.1                    | 26.4  |
| 0.30                 | 25.3           | 23.8                    | 26.9  |
| 0.35                 | 25.9           | 24.3                    | 27.4  |
| 0.40                 | 26.4           | 24.9                    | 27.8  |
| 0.45                 | 26.8           | 25.4                    | 28.3  |
| 0.50                 | 27.3           | 25.9                    | 28.7  |
| 0.55                 | 27.8           | 26.3                    | 29.2  |
| 0.60                 | 28.2           | 26.8                    | 29.7  |
| 0.65                 | 28.7           | 27.2                    | 30.3  |
| 0.70                 | 29.3           | 27.7                    | 30.9  |
| 0.75                 | 29.9           | 28.2                    | 31.5  |
| 0.80                 | 30.5           | 28.7                    | 32.3  |
| 0.85                 | 31.3           | 29.4                    | 33.3  |
| 0.90                 | 32.4           | 30.2                    | 34.6  |
| 0.95                 | 34.1           | 31.5                    | 36.8  |
| 0.99                 | 38.0           | 34.2                    | 41.8  |
| Males                |                |                         |       |
| 0.05                 | 19.0           | 16.2                    | 21.7  |
| 0.10                 | 21.0           | 18.7                    | 23.2  |
| 0.15                 | 22.2           | 20.3                    | 24.1  |
| 0.20                 | 23.1           | 21.4                    | 24.8  |
| 0.25                 | 23.9           | 22.3                    | 25.4  |
| 0.30                 | 24.5           | 23.1                    | 26.0  |
| 0.35                 | 25.1           | 23.8                    | 26.5  |
| 0.40                 | 25.7           | 24.4                    | 27.0  |
| 0.45                 | 26.3           | 25.0                    | 27.5  |
| 0.50                 | 26.8           | 25.5                    | 28.0  |
| 0.55                 | 27.3           | 26.1                    | 28.6  |
| 0.60                 | 27.9           | 26.6                    | 29.1  |
| 0.65                 | 28.4           | 27.1                    | 29.8  |
| 0.70                 | 29.0           | 27.6                    | 30.4  |
| 0.75                 | 29.7           | 28.2                    | 31.2  |
| 0.80                 | 30.5           | 28.8                    | 32.1  |
| 0.85                 | 31.4           | 29.6                    | 33.2  |
| 0.90                 | 32.6           | 30.5                    | 34.7  |
| 0.95                 | 34.6           | 32.0                    | 37.2  |
| 0.99                 | 39.0           | 35.1                    | 42.8  |

<sup>a</sup> Offshore strata 24–26.

**Figure 32**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of Cape Cod female ( $n=140$ ) and male ( $n=165$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 34**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Cape Cod female ( $n=140$ ) and male ( $n=165$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.00                 | 1.0  | 0.5                     | 1.5   |
| 0.08                 | 2.0  | 1.8                     | 2.2   |
| 0.81                 | 3.0  | 2.8                     | 3.2   |
| 1.00                 | 4.0  | 3.6                     | 4.4   |
| 1.00                 | 5.0  | 4.3                     | 5.7   |
| 1.00                 | 6.0  | 5.0                     | 7.0   |
| 1.00                 | 7.0  | 5.7                     | 8.3   |
| 1.00                 | 8.0  | 6.4                     | 9.6   |
| Males                |      |                         |       |
| 0.01                 | 1.0  | 0.5                     | 1.5   |
| 0.16                 | 2.0  | 1.8                     | 2.2   |
| 0.77                 | 3.0  | 2.8                     | 3.2   |
| 0.98                 | 4.0  | 3.6                     | 4.4   |
| 1.00                 | 5.0  | 4.3                     | 5.7   |
| 1.00                 | 6.0  | 5.0                     | 7.0   |
| 1.00                 | 7.0  | 5.7                     | 8.3   |
| 1.00                 | 8.0  | 6.4                     | 9.6   |
| 1.00                 | 9.0  | 7.1                     | 10.9  |
| 1.00                 | 10.0 | 7.8                     | 12.2  |
| 1.00                 | 11.0 | 8.5                     | 13.5  |

<sup>a</sup> Offshore strata 24–26.




**Figure 33**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of Southern New England female ( $n=204$ ) and male ( $n=367$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 35**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Southern New England female ( $n=204$ ) and male ( $n=367$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 18.9           | 14.5                    | 23.3  |
| 0.10                 | 20.6           | 16.9                    | 24.3  |
| 0.15                 | 21.6           | 18.4                    | 24.9  |
| 0.20                 | 22.4           | 19.5                    | 25.4  |
| 0.25                 | 23.1           | 20.4                    | 25.7  |
| 0.30                 | 23.6           | 21.2                    | 26.1  |
| 0.35                 | 24.1           | 21.9                    | 26.4  |
| 0.40                 | 24.6           | 22.5                    | 26.7  |
| 0.45                 | 25.1           | 23.2                    | 27.0  |
| 0.50                 | 25.5           | 23.8                    | 27.3  |
| 0.55                 | 26.0           | 24.4                    | 27.6  |
| 0.60                 | 26.4           | 25.0                    | 27.9  |
| 0.65                 | 26.9           | 25.6                    | 28.3  |
| 0.70                 | 27.4           | 26.2                    | 28.7  |
| 0.75                 | 28.0           | 26.9                    | 29.1  |
| 0.80                 | 28.6           | 27.5                    | 29.7  |
| 0.85                 | 29.4           | 28.3                    | 30.6  |
| 0.90                 | 30.5           | 29.1                    | 31.8  |
| 0.95                 | 32.1           | 30.3                    | 34.0  |
| 0.99                 | 35.8           | 32.6                    | 39.1  |
| Males                |                |                         |       |
| 0.05                 | 11.7           | 6.5                     | 16.9  |
| 0.10                 | 13.7           | 9.2                     | 18.2  |
| 0.15                 | 14.9           | 10.8                    | 19.1  |
| 0.20                 | 15.9           | 12.0                    | 19.7  |
| 0.25                 | 16.6           | 13.0                    | 20.2  |
| 0.30                 | 17.3           | 13.9                    | 20.7  |
| 0.35                 | 17.9           | 14.7                    | 21.1  |
| 0.40                 | 18.5           | 15.5                    | 21.5  |
| 0.45                 | 19.0           | 16.2                    | 21.9  |
| 0.50                 | 19.6           | 16.9                    | 22.3  |
| 0.55                 | 20.1           | 17.5                    | 22.7  |
| 0.60                 | 20.6           | 18.2                    | 23.0  |
| 0.65                 | 21.2           | 19.0                    | 23.5  |
| 0.70                 | 21.8           | 19.7                    | 23.9  |
| 0.75                 | 22.5           | 20.5                    | 24.4  |
| 0.80                 | 23.3           | 21.5                    | 25.1  |
| 0.85                 | 24.2           | 22.6                    | 25.8  |
| 0.90                 | 25.4           | 23.9                    | 26.9  |
| 0.95                 | 27.4           | 25.9                    | 29.0  |
| 0.99                 | 31.8           | 29.3                    | 34.3  |

<sup>a</sup> Offshore strata 5–6, 9–10.




**Figure 34**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of Southern New England female ( $n=204$ ) and male ( $n=367$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 36**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Southern New England female ( $n=204$ ) and male ( $n=367$ ) yellowtail flounder, *Pleuronectes ferrugineus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Age | 95% Confidence interval |       |
|----------------------|-----|-------------------------|-------|
|                      |     | Lower                   | Upper |
| Females              |     |                         |       |
| 0.13                 | 1.0 | 0.6                     | 1.4   |
| 0.74                 | 2.0 | 1.9                     | 2.1   |
| 0.98                 | 3.0 | 2.6                     | 3.4   |
| 1.00                 | 4.0 | 3.3                     | 4.7   |
| 1.00                 | 5.0 | 3.9                     | 6.1   |
| 1.00                 | 6.0 | 4.6                     | 7.4   |
| Males                |     |                         |       |
| 0.00                 | 1.0 | -1.3                    | 3.3   |
| 0.92                 | 2.0 | 1.9                     | 2.1   |
| 1.00                 | 3.0 | 0.7                     | 5.3   |
| 1.00                 | 4.0 | -0.7                    | 8.7   |
| 1.00                 | 5.0 | -2.0                    | 12.0  |
| 1.00                 | 6.0 | -3.4                    | 15.4  |

<sup>a</sup> Offshore strata 5–6, 9–10.

## Winter Flounder: *Pleuronectes americanus*

Winter flounder are widely distributed in the northwest Atlantic from Labrador to Georgia, but abundance is highest from the Gulf of St. Lawrence to Chesapeake Bay. Tagging and meristic studies indicate three stocks of winter flounder in U.S. waters: two inshore stocks north and south of Cape Cod (Lux et al. 1970; Howe and Coates 1975; Pierce and Howe 1977) and an offshore stock occupying Georges Bank (Lux 1973). Spawning occurs from February to May, with peak activity during February and March in southern areas and April farther north (Table 2). Commercial landings of winter flounder from Georges Bank averaged 3,800 metric tons per year during 1980–1984 but have subsequently declined to about 2,300 t annually since then (NEFSC 1991). Combined commercial and recreational catch for inshore stocks has similarly declined in recent years, from about 6,400 t per year during 1980–1982 to 2,700 t annually since 1983 for the northern stock, and from an average of 14,000 t per year during 1980–1985 to 7,800 t from 1986–1989 for the southern stock (NEFSC 1991).




**Figure 35**

Calculated and observed (•) proportion mature at length based on macroscopic observations of whole gonads of Georges Bank female ( $n=204$ ) and male ( $n=174$ ) winter flounder, *Pleuronectes americanus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1989.

Median length at maturity for female and male winter flounder from Georges Bank was 24.9 and 25.6 cm, respectively (Tables 3 and 37, Fig. 35). For inshore

**Table 37**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Georges Bank female ( $n=204$ ) and male ( $n=174$ ) winter flounder, *Pleuronectes americanus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Female               |                |                         |       |
| 0.05                 | 18.2           | 14.8                    | 21.6  |
| 0.10                 | 19.9           | 17.1                    | 22.7  |
| 0.15                 | 20.9           | 18.5                    | 23.4  |
| 0.20                 | 21.7           | 19.5                    | 23.9  |
| 0.25                 | 22.4           | 20.4                    | 24.4  |
| 0.30                 | 23.0           | 21.2                    | 24.8  |
| 0.35                 | 23.5           | 21.8                    | 25.2  |
| 0.40                 | 24.0           | 22.4                    | 25.6  |
| 0.45                 | 24.5           | 23.0                    | 25.9  |
| 0.50                 | 24.9           | 23.5                    | 26.3  |
| 0.55                 | 25.4           | 24.1                    | 26.7  |
| 0.60                 | 25.9           | 24.6                    | 27.1  |
| 0.65                 | 26.3           | 25.1                    | 27.6  |
| 0.70                 | 26.9           | 25.6                    | 28.1  |
| 0.75                 | 27.4           | 26.1                    | 28.8  |
| 0.80                 | 28.1           | 26.7                    | 29.5  |
| 0.85                 | 28.9           | 27.3                    | 30.5  |
| 0.90                 | 30.0           | 28.1                    | 31.8  |
| 0.95                 | 31.7           | 29.3                    | 34.0  |
| 0.99                 | 35.5           | 31.8                    | 39.1  |
| Males                |                |                         |       |
| 0.05                 | 15.9           | 11.1                    | 20.6  |
| 0.10                 | 18.3           | 14.4                    | 22.2  |
| 0.15                 | 19.9           | 16.4                    | 23.3  |
| 0.20                 | 21.0           | 17.9                    | 24.1  |
| 0.25                 | 22.0           | 19.2                    | 24.8  |
| 0.30                 | 22.8           | 20.2                    | 25.4  |
| 0.35                 | 23.6           | 21.2                    | 25.9  |
| 0.40                 | 24.3           | 22.1                    | 26.5  |
| 0.45                 | 25.0           | 22.9                    | 27.0  |
| 0.50                 | 25.6           | 23.7                    | 27.5  |
| 0.55                 | 26.3           | 24.5                    | 28.1  |
| 0.60                 | 27.0           | 25.2                    | 28.7  |
| 0.65                 | 27.7           | 26.0                    | 29.4  |
| 0.70                 | 28.4           | 26.8                    | 30.1  |
| 0.75                 | 29.3           | 27.6                    | 31.0  |
| 0.80                 | 30.2           | 28.4                    | 32.0  |
| 0.85                 | 31.4           | 29.4                    | 33.4  |
| 0.90                 | 32.9           | 30.6                    | 35.2  |
| 0.95                 | 35.4           | 32.4                    | 38.4  |
| 0.99                 | 40.9           | 36.1                    | 45.6  |

<sup>a</sup> Offshore strata 13–22.

stocks, values of  $L_{50}$  were 29.7 cm for females and 27.6 cm for males north of Cape Cod, and 27.6 cm for females and 29.0 cm for males south of Cape Cod (Tables 3, 39, and 41, Figs. 37 and 39). Median age at maturity was 1.9 yr for both sexes on Georges Bank (Table 4, Fig. 36); proportions of mature fish calculated for each age are presented in Table 38. For inshore stocks, values of  $A_{50}$  were 3.5 yr for females and 3.3 yr for females north of Cape Cod, and 3.0 yr for females and 3.3 yr for males south of Cape Cod (Table 4, Figs. 38 and 40); proportions mature calculated for each age are presented in Tables 40 and 42.

Morse (1979)<sup>2</sup> reported values for  $L_{50}$  derived from 1977 data of 26.3 cm for females and 24.8 cm for males, but did not distinguish between inshore stocks and the faster-growing (Lux 1973) Georges Bank winter flounder. Comparisons of maturity analyses from other stud-

ies are complicated by the complex stock structure for the species. Perlmutter (1947) observed that winter flounder in the Long Island Sound region mature at 20–25 cm and 2–3 yr of age, while Kennedy and Steele (1971) reported values for  $L_{50}$  of 25 cm for females and 21 cm for males in Newfoundland waters, with ages at full maturity of 7 yr for females and 6 yr for males, leaving these authors to conclude that maturation was related to size and not age. However, Beacham (1982) presented evidence that maturation of winter flounder in the Scotian Shelf and southern Gulf of St. Lawrence region was highly variable from year to year, and Burton and Idler (1984) found large numbers of nonreproductive individuals in a given year, suggesting that interpretations of maturity data for winter flounder must be conducted with caution.




**Figure 36**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of Georges Bank female ( $n=204$ ) and male ( $n=174$ ) winter flounder, *Pleuronectes americanus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1989.

**Table 38**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of Georges Bank female ( $n=204$ ) and male ( $n=174$ ) winter flounder, *Pleuronectes americanus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.03                 | 1.0  | 0.4                     | 1.6   |
| 0.62                 | 2.0  | 1.9                     | 2.1   |
| 0.99                 | 3.0  | 2.5                     | 3.5   |
| 1.00                 | 4.0  | 3.0                     | 5.0   |
| 1.00                 | 5.0  | 3.5                     | 6.5   |
| 1.00                 | 6.0  | 3.9                     | 8.1   |
| 1.00                 | 7.0  | 4.4                     | 9.6   |
| 1.00                 | 8.0  | 4.9                     | 11.1  |
| 1.00                 | 9.0  | 5.4                     | 12.6  |
| 1.00                 | 10.0 | 5.8                     | 14.2  |
| Males                |      |                         |       |
| 0.10                 | 1.0  | 0.5                     | 1.5   |
| 0.56                 | 2.0  | 1.8                     | 2.2   |
| 0.94                 | 3.0  | 2.6                     | 3.4   |
| 0.99                 | 4.0  | 3.2                     | 4.8   |
| 1.00                 | 5.0  | 3.8                     | 6.2   |
| 1.00                 | 6.0  | 4.4                     | 7.6   |
| 1.00                 | 7.0  | 5.0                     | 9.0   |
| 1.00                 | 8.0  | 5.6                     | 10.4  |
| 1.00                 | 9.0  | 6.2                     | 11.8  |

<sup>a</sup> Offshore strata 13–22.




**Figure 37**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=320$ ) and male ( $n=215$ ) winter flounder, *Pleuronectes americanus*, sampled north of Cape Cod during Massachusetts Division of Marine Fisheries spring bottom trawl surveys, 1985–1989.

**Table 39**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=320$ ) and male ( $n=215$ ) winter flounder, *Pleuronectes americanus*, sampled north of Cape Cod during Massachusetts Division of Marine Fisheries spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 26.5           | 25.3                    | 27.6  |
| 0.10                 | 27.3           | 26.4                    | 28.2  |
| 0.15                 | 27.8           | 27.0                    | 28.6  |
| 0.20                 | 28.2           | 27.5                    | 28.9  |
| 0.25                 | 28.5           | 27.8                    | 29.2  |
| 0.30                 | 28.8           | 28.2                    | 29.4  |
| 0.35                 | 29.0           | 28.5                    | 29.6  |
| 0.40                 | 29.3           | 28.7                    | 29.8  |
| 0.45                 | 29.5           | 29.0                    | 30.1  |
| 0.50                 | 29.7           | 29.2                    | 30.3  |
| 0.55                 | 30.0           | 29.5                    | 30.5  |
| 0.60                 | 30.2           | 29.7                    | 30.7  |
| 0.65                 | 30.4           | 29.9                    | 30.9  |
| 0.70                 | 30.7           | 30.2                    | 31.2  |
| 0.75                 | 31.0           | 30.4                    | 31.5  |
| 0.80                 | 31.3           | 30.7                    | 31.9  |
| 0.85                 | 31.7           | 31.0                    | 32.3  |
| 0.90                 | 32.2           | 31.5                    | 32.9  |
| 0.95                 | 33.0           | 32.1                    | 33.9  |
| 0.99                 | 34.9           | 33.5                    | 36.2  |
| Males                |                |                         |       |
| 0.05                 | 22.8           | 21.2                    | 24.3  |
| 0.10                 | 24.0           | 22.8                    | 25.2  |
| 0.15                 | 24.8           | 23.7                    | 25.8  |
| 0.20                 | 25.3           | 24.4                    | 26.3  |
| 0.25                 | 25.8           | 24.9                    | 26.7  |
| 0.30                 | 26.2           | 25.4                    | 27.0  |
| 0.35                 | 26.6           | 25.8                    | 27.3  |
| 0.40                 | 26.9           | 26.2                    | 27.6  |
| 0.45                 | 27.3           | 26.6                    | 28.0  |
| 0.50                 | 27.6           | 26.9                    | 28.3  |
| 0.55                 | 27.9           | 27.2                    | 28.6  |
| 0.60                 | 28.2           | 27.5                    | 29.0  |
| 0.65                 | 28.6           | 27.9                    | 29.3  |
| 0.70                 | 29.0           | 28.2                    | 29.7  |
| 0.75                 | 29.4           | 28.5                    | 30.2  |
| 0.80                 | 29.8           | 28.9                    | 30.8  |
| 0.85                 | 30.4           | 29.4                    | 31.4  |
| 0.90                 | 31.2           | 30.0                    | 32.4  |
| 0.95                 | 32.4           | 30.9                    | 33.9  |
| 0.99                 | 35.1           | 32.9                    | 37.3  |

<sup>a</sup> Strata 25–36.




**Figure 38**


Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=320$ ) and male ( $n=215$ ) winter flounder, *Pleuronectes americanus*, sampled north of Cape Cod during Massachusetts Division of Marine Fisheries spring bottom trawl surveys, 1985–1989.

**Table 40**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=320$ ) and male ( $n=215$ ) winter flounder, *Pleuronectes americanus*, sampled north of Cape Cod during Massachusetts Division of Marine Fisheries spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>Mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.00                 | 1.0  | 0.5                     | 1.5   |
| 0.01                 | 2.0  | 1.7                     | 2.3   |
| 0.16                 | 3.0  | 2.9                     | 3.1   |
| 0.86                 | 4.0  | 3.9                     | 4.1   |
| 0.99                 | 5.0  | 4.7                     | 5.3   |
| 1.00                 | 6.0  | 5.5                     | 6.5   |
| 1.00                 | 7.0  | 6.3                     | 7.7   |
| 1.00                 | 8.0  | 7.1                     | 8.9   |
| 1.00                 | 9.0  | 7.9                     | 10.1  |
| 1.00                 | 10.0 | 8.7                     | 11.3  |
| 1.00                 | 11.0 | 9.5                     | 12.5  |
| 1.00                 | 12.0 | 10.3                    | 13.7  |
| 1.00                 | 13.0 | 11.1                    | 14.9  |
| Males                |      |                         |       |
| 0.00                 | 1.0  | 0.4                     | 1.6   |
| 0.04                 | 2.0  | 1.6                     | 2.4   |
| 0.34                 | 3.0  | 2.9                     | 3.1   |
| 0.87                 | 4.0  | 3.7                     | 4.3   |
| 0.99                 | 5.0  | 4.4                     | 5.6   |
| 1.00                 | 6.0  | 5.2                     | 6.8   |
| 1.00                 | 7.0  | 5.9                     | 8.1   |
| 1.00                 | 8.0  | 6.6                     | 9.4   |
| 1.00                 | 9.0  | 7.3                     | 10.7  |

<sup>a</sup> Strata 25–36.

**Figure 39**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=398$ ) and male ( $n=301$ ) winter flounder, *Pleuronectes americanus*, sampled south of Cape Cod during Massachusetts Division of Marine Fisheries spring bottom trawl surveys, 1985–1989.

**Table 41**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=398$ ) and male ( $n=301$ ) winter flounder, *Pleuronectes americanus*, sampled south of Cape Cod during Massachusetts Division of Marine Fisheries spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 20.8           | 19.0                    | 22.5  |
| 0.10                 | 22.5           | 21.0                    | 23.9  |
| 0.15                 | 23.6           | 22.3                    | 24.8  |
| 0.20                 | 24.4           | 23.2                    | 25.5  |
| 0.25                 | 25.0           | 24.0                    | 26.1  |
| 0.30                 | 25.6           | 24.7                    | 26.6  |
| 0.35                 | 26.1           | 25.3                    | 27.0  |
| 0.40                 | 26.6           | 25.8                    | 27.5  |
| 0.45                 | 27.1           | 26.3                    | 27.9  |
| 0.50                 | 27.6           | 26.8                    | 28.3  |
| 0.55                 | 28.1           | 27.3                    | 28.8  |
| 0.60                 | 28.5           | 27.8                    | 29.3  |
| 0.65                 | 29.0           | 28.3                    | 29.8  |
| 0.70                 | 29.6           | 28.8                    | 30.3  |
| 0.75                 | 30.1           | 29.3                    | 30.9  |
| 0.80                 | 30.8           | 29.9                    | 31.7  |
| 0.85                 | 31.6           | 30.6                    | 32.6  |
| 0.90                 | 32.7           | 31.5                    | 33.8  |
| 0.95                 | 34.4           | 33.0                    | 35.9  |
| 0.99                 | 38.2           | 36.1                    | 40.4  |
| Males                |                |                         |       |
| 0.05                 | 24.0           | 22.6                    | 25.3  |
| 0.10                 | 25.2           | 24.1                    | 26.3  |
| 0.15                 | 26.0           | 25.1                    | 27.0  |
| 0.20                 | 26.6           | 25.8                    | 27.5  |
| 0.25                 | 27.1           | 26.3                    | 27.9  |
| 0.30                 | 27.5           | 26.8                    | 28.3  |
| 0.35                 | 27.9           | 27.2                    | 28.6  |
| 0.40                 | 28.3           | 27.6                    | 29.0  |
| 0.45                 | 28.6           | 28.0                    | 29.3  |
| 0.50                 | 29.0           | 28.3                    | 29.6  |
| 0.55                 | 29.3           | 28.6                    | 30.0  |
| 0.60                 | 29.7           | 29.0                    | 30.4  |
| 0.65                 | 30.0           | 29.3                    | 30.7  |
| 0.70                 | 30.4           | 29.6                    | 31.2  |
| 0.75                 | 30.8           | 30.0                    | 31.7  |
| 0.80                 | 31.3           | 30.4                    | 32.2  |
| 0.85                 | 31.9           | 30.9                    | 32.9  |
| 0.90                 | 32.7           | 31.5                    | 33.8  |
| 0.95                 | 34.0           | 32.5                    | 35.4  |
| 0.99                 | 36.8           | 34.7                    | 38.8  |

<sup>a</sup> Strata 11–17.




**Figure 40**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=398$ ) and male ( $n=301$ ) winter flounder, *Pleuronectes americanus*, sampled south of Cape Cod during Massachusetts Division of Marine Fisheries spring bottom trawl surveys, 1985–1989.

**Table 42**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=398$ ) and male ( $n=301$ ) winter flounder, *Pleuronectes americanus*, sampled south of Cape Cod during Massachusetts Division of Marine Fisheries spring bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.00                 | 1.0  | 0.5                     | 1.5   |
| 0.06                 | 2.0  | 1.7                     | 2.3   |
| 0.53                 | 3.0  | 2.9                     | 3.1   |
| 0.95                 | 4.0  | 3.8                     | 4.2   |
| 1.00                 | 5.0  | 4.5                     | 5.5   |
| 1.00                 | 6.0  | 5.3                     | 6.7   |
| 1.00                 | 7.0  | 6.1                     | 7.9   |
| 1.00                 | 8.0  | 6.8                     | 9.2   |
| 1.00                 | 9.0  | 7.6                     | 10.4  |
| 1.00                 | 10.0 | 8.4                     | 11.6  |
| Males                |      |                         |       |
| 0.00                 | 1.0  | 0.4                     | 1.6   |
| 0.04                 | 2.0  | 1.7                     | 2.3   |
| 0.32                 | 3.0  | 2.9                     | 3.1   |
| 0.83                 | 4.0  | 3.8                     | 4.2   |
| 0.98                 | 5.0  | 4.5                     | 5.5   |
| 1.00                 | 6.0  | 5.3                     | 6.7   |
| 1.00                 | 7.0  | 6.1                     | 7.9   |
| 1.00                 | 8.0  | 6.8                     | 9.2   |
| 1.00                 | 9.0  | 7.6                     | 10.4  |

<sup>a</sup> Strata 11–17.

## American Plaice: *Hippoglossoides platessoides*

American plaice, or dab, occurs on both sides of the north Atlantic. In the northwest Atlantic, American plaice are distributed along the continental shelf from southern Labrador to Rhode Island with large commercial concentrations existing off the coast of New England in the Gulf of Maine - Georges Bank region (Sullivan 1981). Spawning takes place from February through June with peak spawning occurring in April and May (Table 2). Total commercial landings of American plaice increased from a low of about 3,000 metric tons in the mid-1970's to a peak of more than 15,000 t in 1982; landings subsequently declined to levels similar to the mid-1970's (NEFSC 1991).

Median length at maturity of American plaice sampled from the Gulf of Maine region during spring 1986–1990 was 26.8 cm for females and 22.1 for males (Tables 3 and 43, Fig. 41). Median age at maturity for females and males was 3.6 yr and 3.0 yr, respectively (Table 4, Fig. 42); proportions of mature fish calculated for each age are presented in Table 44.

These values are similar to those reported by Sullivan (1981) for females but not for males. Samples taken in the Gulf of Maine in spring 1980 had  $L_{50}$  and  $A_{50}$  values




**Figure 41**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=714$ ) and male ( $n=633$ ) American plaice, *Hippoglossoides platessoides*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1986–1990.

**Table 43**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=714$ ) and male ( $n=633$ ) American plaice, *Hippoglossoides platessoides*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1986–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 19.8           | 18.4                    | 21.2  |
| 0.10                 | 21.6           | 20.4                    | 22.7  |
| 0.15                 | 22.7           | 21.7                    | 23.7  |
| 0.20                 | 23.5           | 22.6                    | 24.4  |
| 0.25                 | 24.2           | 23.3                    | 25.0  |
| 0.30                 | 24.8           | 24.0                    | 25.5  |
| 0.35                 | 25.3           | 24.6                    | 26.0  |
| 0.40                 | 25.8           | 25.1                    | 26.5  |
| 0.45                 | 26.3           | 25.6                    | 27.0  |
| 0.50                 | 26.8           | 26.1                    | 27.4  |
| 0.55                 | 27.3           | 26.6                    | 27.9  |
| 0.60                 | 27.7           | 27.1                    | 28.4  |
| 0.65                 | 28.3           | 27.6                    | 28.9  |
| 0.70                 | 28.8           | 28.1                    | 29.5  |
| 0.75                 | 29.4           | 28.7                    | 30.1  |
| 0.80                 | 30.1           | 29.3                    | 30.9  |
| 0.85                 | 30.9           | 30.0                    | 31.8  |
| 0.90                 | 32.0           | 31.0                    | 33.0  |
| 0.95                 | 33.8           | 32.6                    | 35.0  |
| 0.99                 | 37.7           | 35.9                    | 39.5  |
| Males                |                |                         |       |
| 0.05                 | 13.3           | 11.7                    | 14.9  |
| 0.10                 | 15.5           | 14.2                    | 16.8  |
| 0.15                 | 16.9           | 15.8                    | 18.0  |
| 0.20                 | 18.0           | 17.0                    | 18.9  |
| 0.25                 | 18.8           | 17.9                    | 19.7  |
| 0.30                 | 19.6           | 18.7                    | 20.4  |
| 0.35                 | 20.2           | 19.5                    | 21.0  |
| 0.40                 | 20.9           | 20.2                    | 21.6  |
| 0.45                 | 21.5           | 20.8                    | 22.2  |
| 0.50                 | 22.1           | 21.4                    | 22.7  |
| 0.55                 | 22.7           | 22.0                    | 23.3  |
| 0.60                 | 23.3           | 22.6                    | 24.0  |
| 0.65                 | 23.9           | 23.3                    | 24.6  |
| 0.70                 | 24.6           | 23.9                    | 25.3  |
| 0.75                 | 25.4           | 24.6                    | 26.1  |
| 0.80                 | 26.2           | 25.4                    | 27.1  |
| 0.85                 | 27.3           | 26.3                    | 28.2  |
| 0.90                 | 28.6           | 27.5                    | 29.7  |
| 0.95                 | 30.9           | 29.5                    | 32.3  |
| 0.99                 | 35.8           | 33.7                    | 37.9  |

<sup>a</sup> Offshore strata 26–30, 36–40.

for females of 27.4 cm and 3.4 yr, and 26.5 cm and 3.4 yr for males, respectively. Morse (1979)<sup>2</sup> calculated a larger  $L_{50}$  value of 33.6 cm for females and a similar value of 27.2 cm for males for American plaice sampled from Cape Hatteras, N.C. to Nova Scotia during 1977. Median length and age at maturity of female American

plaice in the Scotian Shelf area declined from 31.0 cm and 6 yr during 1970–1974 to 30.8 cm and 4.7 yr during 1975–1979. Median values for males similarly declined from 24.8 cm and 4.5 yr, respectively, to 21.9 cm and 3.5 yr, respectively, during the same time periods (Beacham 1983d).




**Figure 42**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=714$ ) and male ( $n=633$ ) American plaice, *Hippoglossoides platessoides*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1986–1990.

**Table 44**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=714$ ) and male ( $n=633$ ) American plaice, *Hippoglossoides platessoides*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1986–1990.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| <b>Females</b>       |      |                         |       |
| 0.00                 | 1.0  | 0.6                     | 1.4   |
| 0.04                 | 2.0  | 1.7                     | 2.3   |
| 0.24                 | 3.0  | 2.9                     | 3.1   |
| 0.72                 | 4.0  | 3.9                     | 4.1   |
| 0.95                 | 5.0  | 4.8                     | 5.2   |
| 0.99                 | 6.0  | 5.6                     | 6.4   |
| 1.00                 | 7.0  | 6.5                     | 7.5   |
| 1.00                 | 8.0  | 7.3                     | 8.7   |
| 1.00                 | 9.0  | 8.2                     | 9.8   |
| 1.00                 | 10.0 | 9.0                     | 11.0  |
| 1.00                 | 11.0 | 9.9                     | 12.1  |
| 1.00                 | 12.0 | 10.7                    | 13.3  |
| <b>Males</b>         |      |                         |       |
| 0.04                 | 1.0  | 0.7                     | 1.3   |
| 0.16                 | 2.0  | 1.8                     | 2.2   |
| 0.50                 | 3.0  | 2.9                     | 3.1   |
| 0.84                 | 4.0  | 3.8                     | 4.2   |
| 0.97                 | 5.0  | 4.7                     | 5.3   |
| 0.99                 | 6.0  | 5.5                     | 6.5   |
| 1.00                 | 7.0  | 6.4                     | 7.6   |
| 1.00                 | 8.0  | 7.2                     | 8.8   |
| 1.00                 | 9.0  | 8.1                     | 9.9   |

<sup>a</sup> Offshore strata 26–30, 36–40.

## Witch Flounder: *Glyptocephalus cynoglossus*

Witch flounder range from Labrador to Cape Hatteras, North Carolina, but are not commercially abundant south of Cape Cod, Massachusetts. The population inhabiting the Gulf of Maine-Georges Bank region is considered to be a unit stock. Spawning in this region is protracted, encompassing a period from early spring to late summer, but peak activity appears to occur during May and June (Table 2). Commercial landings of witch flounder averaged 2,800 metric tons from 1960 to 1981, rose sharply to an average of 5,500 t during 1982–86, and have subsequently declined to 2,100 t in 1989 (NEFSC 1991).

Median length at maturity for female and male witch flounder was 30.4 and 25.3 cm, respectively (Tables 3 and 45, Fig. 43). Median age at maturity was 4.4 yr for females and 3.6 yr for males (Table 4, Fig. 44); proportions of mature fish calculated for each age are presented in Table 46.

Values of  $L_{50}$  above are lower than those obtained for witch flounder in the Gulf of Maine-Georges Bank region by Morse (1979)<sup>2</sup> from 1977 data (34.9 cm for

females and 33.0 cm for males) and by Burnett et al. (1992) for the period 1977–84 (33.5 and 27.6 cm for females and males, respectively). Additionally,  $A_{50}$  for

**Table 45**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=189$ ) and male ( $n=233$ ) witch flounder, *Glyptocephalus cynoglossus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1986–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 22.9           | 18.8                    | 27.0  |
| 0.10                 | 24.8           | 21.5                    | 28.2  |
| 0.15                 | 26.0           | 23.1                    | 28.9  |
| 0.20                 | 26.9           | 24.3                    | 29.5  |
| 0.25                 | 27.6           | 25.2                    | 30.0  |
| 0.30                 | 28.3           | 26.1                    | 30.5  |
| 0.35                 | 28.8           | 26.8                    | 30.9  |
| 0.40                 | 29.4           | 27.5                    | 31.3  |
| 0.45                 | 29.9           | 28.1                    | 31.7  |
| 0.50                 | 30.4           | 28.7                    | 32.2  |
| 0.55                 | 30.9           | 29.3                    | 32.6  |
| 0.60                 | 31.5           | 29.8                    | 33.1  |
| 0.65                 | 32.0           | 30.4                    | 33.7  |
| 0.70                 | 32.6           | 30.9                    | 34.3  |
| 0.75                 | 33.2           | 31.5                    | 35.0  |
| 0.80                 | 34.0           | 32.1                    | 35.9  |
| 0.85                 | 34.9           | 32.7                    | 37.0  |
| 0.90                 | 36.0           | 33.6                    | 38.5  |
| 0.95                 | 37.9           | 34.8                    | 41.1  |
| 0.99                 | 42.1           | 37.4                    | 46.9  |
| Males                |                |                         |       |
| 0.05                 | 17.4           | 12.6                    | 22.2  |
| 0.10                 | 19.4           | 15.4                    | 23.5  |
| 0.15                 | 20.7           | 17.1                    | 24.2  |
| 0.20                 | 21.6           | 18.4                    | 24.8  |
| 0.25                 | 22.4           | 19.4                    | 25.3  |
| 0.30                 | 23.0           | 20.3                    | 25.8  |
| 0.35                 | 23.7           | 21.1                    | 26.2  |
| 0.40                 | 24.2           | 21.9                    | 26.6  |
| 0.45                 | 24.8           | 22.6                    | 26.9  |
| 0.50                 | 25.3           | 23.3                    | 27.3  |
| 0.55                 | 25.9           | 24.0                    | 27.7  |
| 0.60                 | 26.4           | 24.7                    | 28.2  |
| 0.65                 | 27.0           | 25.3                    | 28.7  |
| 0.70                 | 27.6           | 26.0                    | 29.2  |
| 0.75                 | 28.3           | 26.7                    | 29.8  |
| 0.80                 | 29.1           | 27.5                    | 30.6  |
| 0.85                 | 30.0           | 28.3                    | 31.7  |
| 0.90                 | 31.2           | 29.3                    | 33.2  |
| 0.95                 | 33.3           | 30.7                    | 35.8  |
| 0.99                 | 37.7           | 33.6                    | 41.8  |

<sup>a</sup> Offshore strata 22–30, 33–34, 351, 36–40.




**Figure 43**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=189$ ) and male ( $n=233$ ) witch flounder, *Glyptocephalus cynoglossus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1986–1990.

both sexes has been reduced by about a year from 1977–84 values (Burnett et al. 1992).

In general, maturation of witch flounder in the Gulf of Maine-Georges Bank region occurs at a similar size but at an earlier age than slower-growing fish in colder Canadian waters. Beacham (1983e) reported ranges for  $L_{50}$  of 33.0 to 34.3 cm for females and 29.2 to 33.0 cm for males off Nova Scotia, with corresponding ranges for  $A_{50}$  of 7.2 to 8.8 yr and 5.1 to 9.2 yr. Bowering (1976) and Bowering and Brodie (1984) observed that maturation of female witch flounder in Newfoundland waters and the Gulf of St. Lawrence occurred at a greater size and age than in any other region (values of  $L_{50}$  ranged from 40.0 to 50.0 cm off Newfoundland and 39.8 to 45.0 cm in the Gulf of St. Lawrence; associated values of  $A_{50}$  ranged from 8.4 to 10.2 yr and 9.8 to 14.5 yr), but that maturation of males was comparable to that noted by Beacham (1983e). Recently, Bowering (1987, 1989) has related changes in biological parameters, including maturation rates, for witch flounder in the Labrador and Newfoundland regions to commercial exploitation patterns.




Figure 44

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=189$ ) and male ( $n=233$ ) witch flounder, *Glyptocephalus cynoglossus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1986–1990.

Table 46

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=189$ ) and male ( $n=233$ ) witch flounder, *Glyptocephalus cynoglossus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1986–1990.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.00                 | 1.0  | -0.5                    | 2.5   |
| 0.02                 | 2.0  | 0.8                     | 3.2   |
| 0.10                 | 3.0  | 2.2                     | 3.8   |
| 0.36                 | 4.0  | 3.5                     | 4.5   |
| 0.73                 | 5.0  | 4.6                     | 5.4   |
| 0.93                 | 6.0  | 5.4                     | 6.6   |
| 0.98                 | 7.0  | 6.0                     | 8.0   |
| 1.00                 | 8.0  | 6.6                     | 9.4   |
| 1.00                 | 9.0  | 7.2                     | 10.8  |
| 1.00                 | 10.0 | 7.8                     | 12.2  |
| 1.00                 | 11.0 | 8.4                     | 13.6  |
| 1.00                 | 12.0 | 9.0                     | 15.0  |
| 1.00                 | 13.0 | 9.6                     | 16.4  |
| 1.00                 | 14.0 | 10.2                    | 17.8  |
| 1.00                 | 15.0 | 10.8                    | 19.2  |
| 1.00                 | 16.0 | 11.4                    | 20.6  |
| 1.00                 | 17.0 | 12.0                    | 22.0  |
| 1.00                 | 18.0 | 12.6                    | 23.4  |
| 1.00                 | 19.0 | 13.2                    | 24.8  |
| 1.00                 | 20.0 | 13.8                    | 26.2  |
| 1.00                 | 21.0 | 14.4                    | 27.6  |
| 1.00                 | 22.0 | 15.0                    | 29.0  |
| 1.00                 | 23.0 | 15.6                    | 30.4  |
| 1.00                 | 24.0 | 16.2                    | 31.8  |
| Males                |      |                         |       |
| 0.00                 | 1.0  | -0.5                    | 2.5   |
| 0.02                 | 2.0  | 1.0                     | 3.0   |
| 0.19                 | 3.0  | 2.5                     | 3.5   |
| 0.71                 | 4.0  | 3.8                     | 4.2   |
| 0.96                 | 5.0  | 4.5                     | 5.5   |
| 1.00                 | 6.0  | 5.0                     | 7.0   |
| 1.00                 | 7.0  | 5.6                     | 8.4   |
| 1.00                 | 8.0  | 6.1                     | 9.9   |
| 1.00                 | 9.0  | 6.6                     | 11.4  |
| 1.00                 | 10.0 | 7.2                     | 12.8  |
| 1.00                 | 11.0 | 7.7                     | 14.3  |
| 1.00                 | 12.0 | 8.2                     | 15.8  |
| 1.00                 | 13.0 | 8.7                     | 17.3  |
| 1.00                 | 14.0 | 9.3                     | 18.7  |
| 1.00                 | 15.0 | 9.8                     | 20.2  |
| 1.00                 | 16.0 | 10.3                    | 21.7  |
| 1.00                 | 17.0 | 10.9                    | 23.1  |
| 1.00                 | 18.0 | 11.4                    | 24.6  |
| 1.00                 | 19.0 | 11.9                    | 26.1  |
| 1.00                 | 20.0 | 12.4                    | 27.6  |
| 1.00                 | 21.0 | 13.0                    | 29.0  |

<sup>a</sup> Offshore strata 22–30, 33–34, 351, 36–40.

## Summer Flounder: *Paralichthys dentatus*

Summer flounder occur from the southern Gulf of Maine to South Carolina but are most common between Cape Cod and Cape Hatteras, where a single stock is thought to exist (Fogarty et al. 1983). Spawning takes place during an autumn migration from coastal embayments to the outer continental shelf with most activity occurring in October and November (Table 2). Commercial landings of summer flounder increased sharply from less than 6,000 metric tons annually prior to 1974 to about 12,000 t per year during the 1980's. An equivalent amount of recreational landings was also taken during the same period (NEFSC 1991).

Median length at maturity for female and male summer flounder was 28.0 and 24.9 cm, respectively (Tables 3 and 47, Fig. 45). These sizes occur at about age 2.5 for females and age 2 for males according to the growth functions presented by Penttila et al. (1989) for the entire northwest Atlantic.

Values of  $L_{50}$  above are comparable to those for both sexes reported by Morse (1979)<sup>2</sup> for the year 1977.




**Figure 45**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=378$ ) and male ( $n=497$ ) summer flounder, *Paralichthys dentatus*, sampled during Northeast Fisheries Science Center autumn bottom trawl surveys, 1985–1989.

Gillikin and Holland (1983), however, suggested that female summer flounder collected off the coast of North Carolina in late 1975 and early 1976 first matured at

**Table 47**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=378$ ) and male ( $n=497$ ) summer flounder, *Paralichthys dentatus*, sampled during Northeast Fisheries Science Center autumn bottom trawl surveys<sup>a</sup>, 1985–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 18.4           | 15.4                    | 21.3  |
| 0.10                 | 20.8           | 18.4                    | 23.3  |
| 0.15                 | 22.3           | 20.2                    | 24.5  |
| 0.20                 | 23.5           | 21.5                    | 25.4  |
| 0.25                 | 24.4           | 22.6                    | 26.2  |
| 0.30                 | 25.2           | 23.5                    | 26.9  |
| 0.35                 | 26.0           | 24.4                    | 27.5  |
| 0.40                 | 26.7           | 25.2                    | 28.1  |
| 0.45                 | 27.3           | 25.9                    | 28.7  |
| 0.50                 | 28.0           | 26.6                    | 29.3  |
| 0.55                 | 28.6           | 27.4                    | 29.9  |
| 0.60                 | 29.3           | 28.1                    | 30.6  |
| 0.65                 | 30.0           | 28.8                    | 31.2  |
| 0.70                 | 30.8           | 29.5                    | 32.0  |
| 0.75                 | 31.6           | 30.3                    | 32.8  |
| 0.80                 | 32.5           | 31.2                    | 33.8  |
| 0.85                 | 33.6           | 32.2                    | 35.1  |
| 0.90                 | 35.2           | 33.6                    | 36.8  |
| 0.95                 | 37.6           | 35.6                    | 39.6  |
| 0.99                 | 43.0           | 40.0                    | 46.0  |
| Males                |                |                         |       |
| 0.05                 | 15.3           | 12.7                    | 17.9  |
| 0.10                 | 17.7           | 15.5                    | 19.9  |
| 0.15                 | 19.2           | 17.3                    | 21.2  |
| 0.20                 | 20.4           | 18.6                    | 22.1  |
| 0.25                 | 21.3           | 19.7                    | 22.9  |
| 0.30                 | 22.1           | 20.6                    | 23.6  |
| 0.35                 | 22.9           | 21.5                    | 24.2  |
| 0.40                 | 23.6           | 22.3                    | 24.8  |
| 0.45                 | 24.2           | 23.0                    | 25.4  |
| 0.50                 | 24.9           | 23.8                    | 26.0  |
| 0.55                 | 25.5           | 24.5                    | 26.6  |
| 0.60                 | 26.2           | 25.2                    | 27.2  |
| 0.65                 | 26.9           | 25.9                    | 27.9  |
| 0.70                 | 27.6           | 26.7                    | 28.6  |
| 0.75                 | 28.5           | 27.5                    | 29.4  |
| 0.80                 | 29.4           | 28.4                    | 30.4  |
| 0.85                 | 30.5           | 29.5                    | 31.6  |
| 0.90                 | 32.0           | 30.8                    | 33.2  |
| 0.95                 | 34.5           | 33.0                    | 36.0  |
| 0.99                 | 39.9           | 37.4                    | 42.3  |

<sup>a</sup> Inshore strata 1–41, offshore strata 1–12, 45–46, 52, 55–76.

age 3 with full maturation occurring after age 4. Their study also indicated that some males matured at age 2, but the majority deferred spawning until age 3 at a median length of 33.1 cm. In a more comprehensive study, Morse (1981) reported an increase in  $L_{50}$  for males from 25.0 to 27.3 cm between 1974–75 and 1978–79. However, there appeared to be no trend in female

$L_{50}$  estimates which fluctuated between 30.2 and 33.3 cm during the same period.

However, a revision to ageing convention (Smith et al. 1981; Almeida et al. 1992) has resulted in median lengths being attained a year earlier than reported above. This interpretation has been supported by recent growth studies by Able et al. (1990) and Szedlmayer et al. (1992).

## Windowpane: *Scophthalmus aquosus*

Windowpane are distributed from the Gulf of St. Lawrence to South Carolina, but are commercially abundant only in the Georges Bank-Southern New England region. No information relative to stock structure exists, but differences in growth rates suggest that fish from Georges Bank and Southern New England comprise separate stocks (Thorpe 1991). Spawning occurs from June to October on Georges Bank, with peak activity during July and August; in Southern New England, spawning takes place from May to November with a peak in September (Table 2). Commercial landings of windowpane have averaged 1,100 metric tons from Georges Bank and 600 t from Southern New England during 1976–1989 (Thorpe 1991).

Median length at maturity for female and male windowpane from Georges Bank was 22.5 and 22.2 cm, respectively, and 21.2 cm for females and 21.5 cm for males from the Southern New England region (Tables 3, 48 and 49; Figs. 46 and 47). Applying the growth function of Moore (1947) to these values of  $L_{50}$  results in estimates of median age at maturity slightly above

age 3; this is consistent with the observation by Moore (1947) that windowpane attain maturity during the third or fourth year.




**Figure 46**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=343$ ) and male ( $n=199$ ) windowpane, *Scophthalmus aquosus*, from Georges Bank sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 48**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=343$ ) and male ( $n=199$ ) windowpane, *Scophthalmus aquosus*, from Georges Bank sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 18.2           | 14.9                    | 21.4  |
| 0.10                 | 19.3           | 16.4                    | 22.1  |
| 0.15                 | 19.9           | 17.4                    | 22.5  |
| 0.20                 | 20.5           | 18.1                    | 22.8  |
| 0.25                 | 20.9           | 18.7                    | 23.1  |
| 0.30                 | 21.2           | 19.2                    | 23.3  |
| 0.35                 | 21.6           | 19.6                    | 23.5  |
| 0.40                 | 21.9           | 20.0                    | 23.8  |
| 0.45                 | 22.2           | 20.4                    | 24.0  |
| 0.50                 | 22.5           | 20.8                    | 24.2  |
| 0.55                 | 22.8           | 21.2                    | 24.4  |
| 0.60                 | 23.1           | 21.6                    | 24.6  |
| 0.65                 | 23.4           | 22.0                    | 24.9  |
| 0.70                 | 23.7           | 22.4                    | 25.1  |
| 0.75                 | 24.1           | 22.8                    | 25.4  |
| 0.80                 | 24.5           | 23.3                    | 25.8  |
| 0.85                 | 25.1           | 23.8                    | 26.3  |
| 0.90                 | 25.7           | 24.4                    | 27.0  |
| 0.95                 | 26.8           | 25.4                    | 28.3  |
| 0.99                 | 29.3           | 27.0                    | 31.5  |
| Males                |                |                         |       |
| 0.05                 | 18.1           | 15.4                    | 20.7  |
| 0.10                 | 19.1           | 16.8                    | 21.4  |
| 0.15                 | 19.8           | 17.7                    | 21.8  |
| 0.20                 | 20.3           | 18.4                    | 22.1  |
| 0.25                 | 20.7           | 19.0                    | 22.4  |
| 0.30                 | 21.0           | 19.4                    | 22.6  |
| 0.35                 | 21.4           | 19.9                    | 22.9  |
| 0.40                 | 21.7           | 20.3                    | 23.1  |
| 0.45                 | 22.0           | 20.6                    | 23.3  |
| 0.50                 | 22.2           | 21.0                    | 23.5  |
| 0.55                 | 22.5           | 21.4                    | 23.7  |
| 0.60                 | 22.8           | 21.7                    | 23.9  |
| 0.65                 | 23.1           | 22.1                    | 24.2  |
| 0.70                 | 23.4           | 22.5                    | 24.4  |
| 0.75                 | 23.8           | 22.9                    | 24.7  |
| 0.80                 | 24.2           | 23.3                    | 25.1  |
| 0.85                 | 24.7           | 23.8                    | 25.6  |
| 0.90                 | 25.4           | 24.3                    | 26.4  |
| 0.95                 | 26.4           | 25.2                    | 27.7  |
| 0.99                 | 28.8           | 26.7                    | 30.8  |

<sup>a</sup> Offshore strata 13–21.




**Figure 47**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=143$ ) and male ( $n=133$ ) windowpane, *Scophthalmus aquosus*, from Southern New England grounds sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1985–1990.

**Table 49**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=143$ ) and male ( $n=133$ ) windowpane, *Scophthalmus aquosus*, from Southern New England grounds sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1985–1990.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 20.5           | 19.5                    | 21.4  |
| 0.10                 | 20.6           | 19.9                    | 21.3  |
| 0.15                 | 20.8           | 20.2                    | 21.3  |
| 0.20                 | 20.8           | 20.4                    | 21.3  |
| 0.25                 | 20.9           | 20.5                    | 21.3  |
| 0.30                 | 21.0           | 20.6                    | 21.4  |
| 0.35                 | 21.0           | 20.6                    | 21.4  |
| 0.40                 | 21.1           | 20.7                    | 21.5  |
| 0.45                 | 21.1           | 20.7                    | 21.6  |
| 0.50                 | 21.2           | 20.7                    | 21.7  |
| 0.55                 | 21.2           | 20.7                    | 21.8  |
| 0.60                 | 21.3           | 20.7                    | 21.9  |
| 0.65                 | 21.3           | 20.7                    | 22.0  |
| 0.70                 | 21.4           | 20.6                    | 22.1  |
| 0.75                 | 21.4           | 20.6                    | 22.2  |
| 0.80                 | 21.5           | 20.6                    | 22.4  |
| 0.85                 | 21.6           | 20.5                    | 22.6  |
| 0.90                 | 21.7           | 20.5                    | 22.9  |
| 0.95                 | 21.9           | 20.4                    | 23.4  |
| 0.99                 | 22.3           | 20.2                    | 24.4  |
| Males                |                |                         |       |
| 0.05                 | 18.3           | 16.3                    | 20.4  |
| 0.10                 | 19.1           | 17.4                    | 20.9  |
| 0.15                 | 19.6           | 18.1                    | 21.2  |
| 0.20                 | 20.0           | 18.6                    | 21.4  |
| 0.25                 | 20.3           | 19.0                    | 21.6  |
| 0.30                 | 20.6           | 19.4                    | 21.8  |
| 0.35                 | 20.9           | 19.7                    | 22.0  |
| 0.40                 | 21.1           | 20.0                    | 22.2  |
| 0.45                 | 21.3           | 20.3                    | 22.3  |
| 0.50                 | 21.5           | 20.6                    | 22.5  |
| 0.55                 | 21.8           | 20.8                    | 22.7  |
| 0.60                 | 22.0           | 21.1                    | 22.9  |
| 0.65                 | 22.2           | 21.3                    | 23.1  |
| 0.70                 | 22.5           | 21.6                    | 23.3  |
| 0.75                 | 22.7           | 21.9                    | 23.6  |
| 0.80                 | 23.1           | 22.2                    | 24.0  |
| 0.85                 | 23.4           | 22.5                    | 24.4  |
| 0.90                 | 24.0           | 22.9                    | 25.0  |
| 0.95                 | 24.8           | 23.5                    | 26.1  |
| 0.99                 | 26.6           | 24.6                    | 28.6  |

<sup>a</sup> Offshore strata 5–12

## Atlantic Herring: *Clupea harengus*

Atlantic herring are widely distributed in the northwest Atlantic from Labrador to Cape Hatteras, North Carolina. Spawning in the Gulf of Maine occurs from late August to December, with peak activity during September and October (Table 2). Historically, important commercial fisheries in U.S. waters have occurred in the coastal areas of the Gulf of Maine and on Georges Bank. In the former region, the fishery targets juvenile herring; in the latter region, no fishery has been prosecuted since the total collapse of the stock in the mid 1970's. Commercial landings of herring averaged 57,000 metric tons per year during 1942–1963, declined to 31,000 t annually from 1964–1975, increased to 60,000 t per year during the period 1976–1981, and have fluctuated around an average of about 33,000 t per year since then (NEFC 1990).<sup>6</sup>

Median length at maturity for female and male Atlantic herring was 25.4 and 25.3 cm (TL), respectively (Tables 3 and 50, Fig. 48). Median age at maturity was 3.0 yr for females and 2.9 yr for males (Table 4, Fig. 49);

proportions of mature fish calculated for each age are presented in Table 51.




**Figure 48**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=359$ ) and male ( $n=307$ ) Atlantic herring, *Clupea harengus*, sampled during Northeast Fisheries Science Center autumn bottom trawl surveys, 1987–1989.

**Table 50**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=359$ ) and male ( $n=307$ ) Atlantic herring, *Clupea harengus*, sampled during Northeast Fisheries Science Center autumn bottom trawl surveys<sup>a</sup>, 1987–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 22.8           | 21.8                    | 23.7  |
| 0.10                 | 23.4           | 22.6                    | 24.2  |
| 0.15                 | 23.8           | 23.1                    | 24.5  |
| 0.20                 | 24.1           | 23.5                    | 24.8  |
| 0.25                 | 24.4           | 23.8                    | 25.0  |
| 0.30                 | 24.6           | 24.1                    | 25.1  |
| 0.35                 | 24.8           | 24.3                    | 25.3  |
| 0.40                 | 25.0           | 24.6                    | 25.5  |
| 0.45                 | 25.2           | 24.8                    | 25.6  |
| 0.50                 | 25.4           | 25.0                    | 25.8  |
| 0.55                 | 25.6           | 25.2                    | 25.9  |
| 0.60                 | 25.7           | 25.4                    | 26.1  |
| 0.65                 | 25.9           | 25.6                    | 26.3  |
| 0.70                 | 26.1           | 25.8                    | 26.5  |
| 0.75                 | 26.4           | 26.0                    | 26.7  |
| 0.80                 | 26.6           | 26.2                    | 27.0  |
| 0.85                 | 26.9           | 26.5                    | 27.3  |
| 0.90                 | 27.3           | 26.9                    | 27.8  |
| 0.95                 | 28.0           | 27.4                    | 28.6  |
| 0.99                 | 29.5           | 28.5                    | 30.4  |
| Males                |                |                         |       |
| 0.05                 | 23.5           | 22.8                    | 24.2  |
| 0.10                 | 24.0           | 23.4                    | 24.5  |
| 0.15                 | 24.2           | 23.7                    | 24.8  |
| 0.20                 | 24.5           | 24.0                    | 24.9  |
| 0.25                 | 24.6           | 24.2                    | 25.1  |
| 0.30                 | 24.8           | 24.4                    | 25.2  |
| 0.35                 | 24.9           | 24.6                    | 25.3  |
| 0.40                 | 25.1           | 24.7                    | 25.4  |
| 0.45                 | 25.2           | 24.9                    | 25.5  |
| 0.50                 | 25.3           | 25.0                    | 25.6  |
| 0.55                 | 25.5           | 25.2                    | 25.8  |
| 0.60                 | 25.6           | 25.3                    | 25.9  |
| 0.65                 | 25.7           | 25.4                    | 26.0  |
| 0.70                 | 25.9           | 25.6                    | 26.2  |
| 0.75                 | 26.0           | 25.7                    | 26.3  |
| 0.80                 | 26.2           | 25.9                    | 26.5  |
| 0.85                 | 26.4           | 26.0                    | 26.8  |
| 0.90                 | 26.7           | 26.3                    | 27.1  |
| 0.95                 | 27.2           | 26.6                    | 27.7  |
| 0.99                 | 28.2           | 27.4                    | 29.0  |

<sup>a</sup> Offshore strata 1–12, 24–30, 33–34, 351, 36–40, 61–76.

Values of  $L_{50}$  and  $A_{50}$  above are comparable to those reported for Nova Scotia herring for the period 1966–75, during which  $L_{50}$  for both sexes combined varied from 25.3 to 27.4 cm (Sinclair et al. 1982), and  $A_{50}$  varied between 2.9 and 3.6 yr for females and 2.8 and 3.6 yr for males (Sinclair et al. 1979). Maturity at age (Table 51) is similar to that obtained for Gulf of Maine herring from monthly commercial sampling during 1985–89 (0.39 of age 3 fish, 0.98 of age 4 fish, 1.00 age 5+; NEFC 1990)<sup>6</sup>. However, Sinclair et al. (1982) noted

a positive relationship between juvenile growth rates and  $L_{50}$ , and Winters (1976) reported decreases in  $A_{50}$  with decreasing adult biomass for herring in the Gulf of St. Lawrence, suggesting that density-dependent processes may influence maturation rates of Atlantic herring.




**Figure 49**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=359$ ) and male ( $n=307$ ) Atlantic herring, *Clupea harengus*, sampled during Northeast Fisheries Science Center autumn bottom trawl surveys, 1987–1989.

<sup>6</sup> NEFC. 1990. Report of the eleventh NEFC Stock Assessment Workshop Fall 1990. NEFC Ref. Doc., 90-09: 58–65.

**Table 51**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=359$ ) and male ( $n=307$ ) Atlantic herring, *Clupea harengus*, sampled during Northeast Fisheries Science Center autumn bottom trawl surveys<sup>a</sup>, 1987–1989.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.00                 | 1.0  | 0.1                     | 1.9   |
| 0.01                 | 2.0  | 1.6                     | 2.4   |
| 0.48                 | 3.0  | 2.9                     | 3.1   |
| 0.99                 | 4.0  | 3.6                     | 4.4   |
| 1.00                 | 5.0  | 4.2                     | 5.8   |
| 1.00                 | 6.0  | 4.8                     | 7.2   |
| 1.00                 | 7.0  | 5.3                     | 8.7   |
| 1.00                 | 8.0  | 5.9                     | 10.1  |
| 1.00                 | 9.0  | 6.5                     | 11.5  |
| 1.00                 | 10.0 | 7.1                     | 12.9  |
| 1.00                 | 11.0 | 7.7                     | 14.3  |
| Males                |      |                         |       |
| 0.00                 | 1.0  | 0.2                     | 1.8   |
| 0.03                 | 2.0  | 1.6                     | 2.4   |
| 0.58                 | 3.0  | 2.9                     | 3.1   |
| 0.99                 | 4.0  | 3.6                     | 4.4   |
| 1.00                 | 5.0  | 4.3                     | 5.7   |
| 1.00                 | 6.0  | 4.9                     | 7.1   |
| 1.00                 | 7.0  | 5.6                     | 8.4   |
| 1.00                 | 8.0  | 6.2                     | 9.8   |
| 1.00                 | 9.0  | 6.8                     | 11.2  |

<sup>a</sup> Offshore strata 1–12, 24–30, 33–34, 351, 36–40, 61–76.

### Atlantic Mackerel: *Scomber scombrus*

The range of the migratory Atlantic mackerel extends from North Carolina to Newfoundland (Sette 1950). Although considered to be a single stock there are two spawning groups: the southern contingent spawns in the Mid-Atlantic Bight from April to July (Table 2) and the northern contingent spawns in the Gulf of St. Lawrence during June and July (NEFSC 1991). Total landings of Atlantic mackerel peaked at about 400,000 metric tons in 1973, then declined to about 30,000 t in the late 1970's, and have recently increased to about 74,000 t in 1989 (NEFSC 1991).

Median length at maturity of female and male Atlantic mackerel was 25.7 cm and 26.0 cm, respectively (Tables 3 and 52, Fig. 50) for samples collected during spring 1987–1989. Median age at maturity was 1.9 yr for both females and males (Table 4, Fig. 51); proportions of mature fish calculated for each age are presented in Table 53.

Mackerel sampled in the Newfoundland area from June–September during 1970–1973 had higher values




**Figure 50**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=697$ ) and male ( $n=770$ ) mackerel, *Scomber scombrus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1987–1989.

for  $L_{50}$  of 34 cm and 35 cm for females and males, respectively (Moores et al. 1975). MacKay (1967) reported that first spawning for mackerel occurred at age 2 and at lengths greater than 30 cm for Atlantic mack-

**Table 52**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=697$ ) and male ( $n=770$ ) mackerel, *Scomber scombrus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1987–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Females              |                |                         |       |
| 0.05                 | 21.1           | 20.0                    | 22.2  |
| 0.10                 | 22.3           | 21.3                    | 23.2  |
| 0.15                 | 23.0           | 22.2                    | 23.8  |
| 0.20                 | 23.5           | 22.8                    | 24.3  |
| 0.25                 | 24.0           | 23.3                    | 24.7  |
| 0.30                 | 24.4           | 23.7                    | 25.0  |
| 0.35                 | 24.7           | 24.1                    | 25.4  |
| 0.40                 | 25.1           | 24.5                    | 25.7  |
| 0.45                 | 25.4           | 24.8                    | 26.0  |
| 0.50                 | 25.7           | 25.1                    | 26.3  |
| 0.55                 | 26.0           | 25.5                    | 26.6  |
| 0.60                 | 26.3           | 25.8                    | 26.9  |
| 0.65                 | 26.7           | 26.1                    | 27.2  |
| 0.70                 | 27.0           | 26.5                    | 27.6  |
| 0.75                 | 27.4           | 26.8                    | 28.0  |
| 0.80                 | 27.9           | 27.3                    | 28.5  |
| 0.85                 | 28.4           | 27.8                    | 29.1  |
| 0.90                 | 29.1           | 28.4                    | 29.9  |
| 0.95                 | 30.3           | 29.4                    | 31.2  |
| 0.99                 | 32.9           | 31.6                    | 34.2  |
| Males                |                |                         |       |
| 0.05                 | 21.5           | 20.5                    | 22.5  |
| 0.10                 | 22.6           | 21.8                    | 23.5  |
| 0.15                 | 23.3           | 22.6                    | 24.1  |
| 0.20                 | 23.9           | 23.2                    | 24.5  |
| 0.25                 | 24.3           | 23.7                    | 24.9  |
| 0.30                 | 24.7           | 24.1                    | 25.3  |
| 0.35                 | 25.0           | 24.5                    | 25.6  |
| 0.40                 | 25.4           | 24.8                    | 25.9  |
| 0.45                 | 25.7           | 25.2                    | 26.2  |
| 0.50                 | 26.0           | 25.5                    | 26.4  |
| 0.55                 | 26.3           | 25.8                    | 26.7  |
| 0.60                 | 26.6           | 26.1                    | 27.0  |
| 0.65                 | 26.9           | 26.5                    | 27.4  |
| 0.70                 | 27.3           | 26.8                    | 27.7  |
| 0.75                 | 27.6           | 27.2                    | 28.1  |
| 0.80                 | 28.1           | 27.6                    | 28.6  |
| 0.85                 | 28.6           | 28.1                    | 29.2  |
| 0.90                 | 29.3           | 28.7                    | 29.9  |
| 0.95                 | 30.4           | 29.7                    | 31.2  |
| 0.99                 | 33.0           | 31.8                    | 34.1  |

<sup>a</sup> Inshore strata 1–81, offshore strata 1–34, 351, 36–76.

erel sampled during May–July 1965–1966, from locations in the Gulf of St. Lawrence and along the coast of Nova Scotia and Massachusetts. The differences in median maturity between these studies and that reported above may be due to the slower growth rate of large year classes that may delay spawning from one to three years (Mackay 1973; Overholtz 1989).




**Figure 51**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=697$ ) and male ( $n=770$ ) Atlantic mackerel, *Scomber scombrus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1987–1989.

**Table 53**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=697$ ) and male ( $n=770$ ) Atlantic mackerel, *Scomber scombrus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1987–1989.

| Proportion<br>mature | Age  | 95% Confidence interval |       |
|----------------------|------|-------------------------|-------|
|                      |      | Lower                   | Upper |
| Females              |      |                         |       |
| 0.02                 | 1.0  | 0.7                     | 1.3   |
| 0.63                 | 2.0  | 1.9                     | 2.1   |
| 0.99                 | 3.0  | 2.7                     | 3.3   |
| 1.00                 | 4.0  | 3.4                     | 4.6   |
| 1.00                 | 5.0  | 4.0                     | 6.0   |
| 1.00                 | 6.0  | 4.7                     | 7.3   |
| 1.00                 | 7.0  | 5.4                     | 8.6   |
| 1.00                 | 8.0  | 6.1                     | 9.9   |
| 1.00                 | 9.0  | 6.8                     | 11.2  |
| 1.00                 | 10.0 | 7.5                     | 12.5  |
| 1.00                 | 11.0 | 8.2                     | 13.8  |
| 1.00                 | 12.0 | 8.8                     | 15.2  |
| 1.00                 | 13.0 | 9.5                     | 16.5  |
| 1.00                 | 14.0 | 10.2                    | 17.8  |
| 1.00                 | 15.0 | 10.9                    | 19.1  |
| 1.00                 | 16.0 | 11.6                    | 20.4  |
| Males                |      |                         |       |
| 0.05                 | 1.0  | 0.8                     | 1.2   |
| 0.58                 | 2.0  | 1.9                     | 2.1   |
| 0.97                 | 3.0  | 2.7                     | 3.3   |
| 1.00                 | 4.0  | 3.5                     | 4.5   |
| 1.00                 | 5.0  | 4.3                     | 5.7   |
| 1.00                 | 6.0  | 5.1                     | 6.9   |
| 1.00                 | 7.0  | 5.9                     | 8.1   |
| 1.00                 | 8.0  | 6.6                     | 9.4   |
| 1.00                 | 9.0  | 7.4                     | 10.6  |
| 1.00                 | 10.0 | 8.2                     | 11.8  |
| 1.00                 | 11.0 | 9.0                     | 13.0  |
| 1.00                 | 12.0 | 9.7                     | 14.3  |
| 1.00                 | 13.0 | 10.5                    | 15.5  |
| 1.00                 | 14.0 | 11.3                    | 16.7  |
| 1.00                 | 15.0 | 12.1                    | 17.9  |
| 1.00                 | 16.0 | 12.8                    | 19.2  |

<sup>a</sup> Inshore strata 1–81, offshore strata 1–34, 351, 36–76.

### Butterfish: *Peprilus triacanthus*

Butterfish are distributed along the Atlantic coast of North America from Newfoundland to Florida, but are most common between Southern New England and Cape Hatteras where a single stock is thought to exist (Murawski and Waring 1979). Spawning takes place in late spring and summer with most activity occurring in June and July (Table 2). Total commercial landings fluctuated between 4,000 and 20,000 metric tons during the 1960's and 1970's, but remained relatively stable at about 5,000 t between 1977 and 1981; landings have since declined to about 2,000 t in 1988 (NEFSC 1991).

Median length at maturity for female and male butterfish was 12.0 and 11.4 cm, respectively (Tables 3 and 54, Fig. 52). Median age at maturity was 0.9 yr for both sexes (Table 4, Fig. 53); proportions of mature fish calculated for each age are presented in Table 55.

Values of  $L_{50}$  above are comparable to those for both sexes reported by Morse (1979)<sup>2</sup> for the year 1977. DuPaul and McEachran (1973) also noted that maturation of butterfish collected from Chesapeake Bay commenced at about age 1 and that almost all two-year-old

fish were sexually mature. In an earlier work, however, Horn (1970) suggested that maturation may have occurred in fish as small as 12 cm but was not common until about 14 cm.




**Figure 52**

Calculated and observed (·) proportion mature at length based on macroscopic observations of whole gonads of female ( $n=333$ ) and male ( $n=341$ ) butterfish, *Peprilus triacanthus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1986–1989.

**Table 54**

Calculated proportion mature at length with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=333$ ) and male ( $n=341$ ) butterfish, *Peprilus triacanthus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1986–1989.

| Proportion<br>mature | Length<br>(cm) | 95% Confidence interval |       |
|----------------------|----------------|-------------------------|-------|
|                      |                | Lower                   | Upper |
| Female               |                |                         |       |
| 0.05                 | 9.5            | 8.7                     | 10.3  |
| 0.10                 | 10.1           | 9.5                     | 10.8  |
| 0.15                 | 10.5           | 9.9                     | 11.1  |
| 0.20                 | 10.8           | 10.3                    | 11.3  |
| 0.25                 | 11.0           | 10.6                    | 11.5  |
| 0.30                 | 11.3           | 10.8                    | 11.7  |
| 0.35                 | 11.4           | 11.0                    | 11.9  |
| 0.40                 | 11.6           | 11.2                    | 12.0  |
| 0.45                 | 11.8           | 11.4                    | 12.2  |
| 0.50                 | 12.0           | 11.6                    | 12.3  |
| 0.55                 | 12.1           | 11.8                    | 12.5  |
| 0.60                 | 12.3           | 11.9                    | 12.6  |
| 0.65                 | 12.5           | 12.1                    | 12.8  |
| 0.70                 | 12.7           | 12.3                    | 13.0  |
| 0.75                 | 12.9           | 12.5                    | 13.2  |
| 0.80                 | 13.1           | 12.7                    | 13.5  |
| 0.85                 | 13.4           | 12.9                    | 13.8  |
| 0.90                 | 13.8           | 13.2                    | 14.3  |
| 0.95                 | 14.4           | 13.7                    | 15.0  |
| 0.99                 | 15.7           | 14.8                    | 16.7  |
| Males                |                |                         |       |
| 0.05                 | 8.4            | 7.5                     | 9.2   |
| 0.10                 | 9.1            | 8.4                     | 9.8   |
| 0.15                 | 9.6            | 9.0                     | 10.2  |
| 0.20                 | 10.0           | 9.4                     | 10.5  |
| 0.25                 | 10.3           | 9.8                     | 10.8  |
| 0.30                 | 10.5           | 10.1                    | 11.0  |
| 0.35                 | 10.8           | 10.3                    | 11.2  |
| 0.40                 | 11.0           | 10.6                    | 11.4  |
| 0.45                 | 11.2           | 10.8                    | 11.6  |
| 0.50                 | 11.4           | 11.0                    | 11.8  |
| 0.55                 | 11.6           | 11.3                    | 12.0  |
| 0.60                 | 11.8           | 11.5                    | 12.2  |
| 0.65                 | 12.0           | 11.7                    | 12.4  |
| 0.70                 | 12.3           | 11.9                    | 12.6  |
| 0.75                 | 12.5           | 12.1                    | 12.9  |
| 0.80                 | 12.8           | 12.4                    | 13.2  |
| 0.85                 | 13.2           | 12.7                    | 13.7  |
| 0.90                 | 13.7           | 13.1                    | 14.2  |
| 0.95                 | 14.4           | 13.8                    | 15.1  |
| 0.99                 | 16.1           | 15.1                    | 17.2  |

<sup>a</sup> Inshore strata 1–46, offshore strata 1–25, 61–76.




**Figure 53**

Calculated and observed (·) proportion mature at age based on macroscopic observations of whole gonads of female ( $n=333$ ) and male ( $n=341$ ) butterfish, *Peprilus triacanthus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys, 1986–1989.

**Table 55**

Calculated proportion mature at age with approximate 95% confidence intervals based on macroscopic observations of whole gonads of female ( $n=333$ ) and male ( $n=341$ ) butterfish, *Peprilus triacanthus*, sampled during Northeast Fisheries Science Center spring bottom trawl surveys<sup>a</sup>, 1986–1989.

| Proportion<br>mature | Age | 95% Confidence interval |       |
|----------------------|-----|-------------------------|-------|
|                      |     | Lower                   | Upper |
| <b>Females</b>       |     |                         |       |
| 0.60                 | 1.0 | 0.9                     | 1.1   |
| 0.97                 | 2.0 | 1.7                     | 2.3   |
| 1.00                 | 3.0 | 2.3                     | 3.7   |
| 1.00                 | 4.0 | 3.0                     | 5.0   |
| <b>Males</b>         |     |                         |       |
| 0.59                 | 1.0 | 0.9                     | 1.1   |
| 0.96                 | 2.0 | 1.7                     | 2.3   |
| 1.00                 | 3.0 | 2.3                     | 3.7   |
| 1.00                 | 4.0 | 3.0                     | 5.0   |

<sup>a</sup> Inshore strata 1–46, offshore strata 1–25, 61–76.

## Discussion

The results presented in this report are intended to describe current maturation patterns for the nineteen included species. However, it is evident from even a cursory comparison of our results with those from earlier periods that, for most species,  $A_{50}$  and, in many cases,  $L_{50}$  values have declined substantially in recent years. While it is often difficult to draw conclusions about changes in maturation from diverse studies which may have employed different model fitting techniques, staging criteria, and seasonal sampling schemes, a comparison between our length-based results and those presented by Morse (1979)<sup>2</sup> is warranted since both studies have drawn on the same Northeast Fisheries Science Center trawl survey data base. Our study expands on the initial work of Morse (1979)<sup>2</sup> to include finer spatial and seasonal resolution, and we have fitted logistic curves rather than employing probit analysis. Given the same set of data, however, both fitting techniques will estimate the inflection point ( $A_{50}$ ,  $L_{50}$ ) with the same degree of accuracy (Ashton 1972). In addition, we have analyzed maturation data collected on cruises conducted only since 1985, with the exception of red hake and redfish, whereas Morse (1979)<sup>2</sup> restricted analyses only to 1977 cruises.

The changes in median size at maturity between 1977 and the mid-to-late 1980's are substantial enough to suggest a real decline rather than a bias resulting from different sampling techniques. Criteria for maturation stages presently in use on NEFSC and MDMF cruises have not changed appreciably since the 1970's and criteria for discriminating between immature and all other mature stages are identical (Burnett et al. 1989). Our initial analyses of the data on an annual basis indicated a gradual decline from the late 1970's through the present. In addition, studies which were conducted in the same general region of the coast prior to the mid-1970's indicate that maturation for most species occurred at older median ages and larger median sizes than those reported by Morse (1979)<sup>2</sup>.

Given these changes, it is clear that current estimates should not be used to represent historical maturation patterns as in the estimation of spawning stock over time in sequential population analyses such as VPA or cohort analysis. Our results are appropriate for use in short-term projections of spawning stock and in spawning stock biomass per recruit analyses (Gabriel et al. 1989) in which the current maturation pattern is expected to represent the short-term equilibrium condition of the stock.

A more comprehensive analysis of the entire data set, with a longer time series, will be required before any definitive conclusions can be drawn about compensatory changes in maturation over time. One such study

(O'Brien 1990) found significant declines in median size and age at maturation in year classes of Georges Bank and Gulf of Maine Atlantic cod, *Gadus morhua*, between 1970 and 1987. Similar results were noted by Bowering (1987) for witch flounder, *Glyptocephalus cynoglossus*, and several reports by Beacham (1983 a, b, and d) indicated long-term changes in maturation for several species on the Scotian Shelf.

To ensure the broadest possible synoptic coverage of the maturity data included in our study, we included observations only from standard bottom trawl survey cruises conducted during spring and autumn. We recognize, however, that such multi-purpose sampling programs often impose some limitations on the utility of certain data. In our case, the spring and autumn coverage of the surveys closely coincided with peak spawning periods for most of the species of flounders, but spring coverage for the winter spawning gadids often occurred 1–2 months after peak spawning. As a result, observations for cod, haddock, pollock, and white hake often contained a large proportion of resting fish whose gonads are most similar superficially to immature specimens. Given this greater potential for misclassification of immature and mature fish, the results obtained for some of the gadiforms may be less reliable than those for pleuronectiforms and other spring or autumn spawners. In particular, we note that median sizes and ages at maturity for cod, haddock, pollock, and white hake obtained from our analyses are considerably lower than previously reported.

Determination of maturation stages based on macroscopic examination of gonads generally involves a degree of subjective judgment on the part of the observer. This difficulty, combined with a high turnover of sea-going personnel from year to year may impart a high degree of variability to the observations unless training is continually monitored as noted by Halliday (1987). Since 1986, the NEFSC Woods Hole Laboratory has conducted a series of workshops during each survey season to monitor and calibrate maturity criteria employed by each scientific party involved in data collection. By examining fresh specimens collectively, members of the various teams are able to coordinate their observations through comparison and discussion of staging criteria. Despite these efforts, however, the inherent subjectivity of the approach remains problematic.

To rectify this potential problem in the future, at-sea maturation observations should be coupled with more objective techniques in order to ascertain the degree of precision and bias of the macroscopic method. Other methods which have demonstrated some success include histological identification of reproductive tissue (Morrison 1990), analysis of plasma parameters (Johnson and Casillas 1991), and comparisons with gonadosomatic indices (O'Brien 1990).

Despite these caveats, our results may be viewed as baseline maturation patterns in relation to current stock conditions (abundance, age structure, sex ratio, etc.). In the future, further monitoring of maturation will be required on a routine basis coincident with proposed management actions designed to increase spawning stock biomass. Any compensatory mechanisms which result in declines in median size and age at maturation accompanying declining spawning stock will likely act in reverse if substantial increases in stock biomass occur over the long-term.

## Acknowledgments

We would like to thank W. Morse, M. Fogarty, and R. Halliday for their critical review of this manuscript. We also appreciate the efforts of all the diligent sea-going personnel whose observations contributed to this analysis.

## Literature cited

- Able, K.W., R.E. Matheson, W.W. Morse, M.P. Fahay, and G. Shepherd. 1990. Patterns of summer flounder, *Paralichthys dentatus*, early life history in the Mid-Atlantic Bight and New Jersey estuaries. Fish. Bull., U.S. 88(1):1-12.
- Alm, G. 1959. Connection between maturity, size, and age in fishes. Inst. Freshwater Res. Drottningholm Rep. 40, 145 p.
- Almeida, F.P. 1987. Stock definition of silver hake in the New England-Middle Atlantic area. North Am. J. Fish. Manage. 7:169-186.
- Almeida, F.P., R.E. Castaneda, R. Jesien, R.C. Greenfield, and John M. Burnett. 1992. Proceedings of the NEFC/ASMFC summer flounder, *Paralichthys dentatus*, ageing workshop, 11-13 June 1990, Northeast Fisheries Center, Woods Hole, MA. NOAA Tech. Mem. NMFS-F/NEC-89, 7 p.
- Ashton, W.D. 1972. The logit transformation with special reference to its uses in bioassay. Griffin and Co., London, 88 p.
- Azarovitz, T.R., J. McGurrian, and R. Seagraves (eds). 1989. Proceedings of a workshop on bottom trawl surveys. Atlantic States Mar. Fish. Comm. Spec. Rep. No. 17, 76 p.
- Battle, H.I. 1951. Contributions to a study of the life history of the hake. Spawning with notes on age determinations. Fish. Res. Board Can. Manuscr. Rep. Biol. Sta. 434, 21 p.
- Beacham, T.D. 1982. Biology and exploitation of winter flounder (*Pseudopleuronectes americanus*) in the Canadian Maritimes area of the northwest Atlantic Ocean. Can. Tech. Rep. Fish. Aquat. Sci. 1113, 33 p.
- 1983a. Variability in median size and age at sexual maturity of Atlantic cod, *Gadus morhua*, on the Scotian Shelf in the northwest Atlantic Ocean. Fish. Bull. U.S. 81(2):303-321.
- 1983b. Variability in size or age at sexual maturity of haddock (*Melanogrammus aeglefinus*) on the Scotian Shelf in the northwest Atlantic Ocean. Can. Tech. Rep. Fish. Aquat. Sci. 1168, iv + 33 p.
- 1983c. Variability in size or age at sexual maturity of white hake, pollock, longfin hake, and silver hake in the Canadian Maritimes region of the northwest Atlantic Ocean. Can. Tech. Rep. Fish. Aquat. Sci. 1157, iv + 43 p.
- 1983d. Variability in size and age at sexual maturity of American plaice and yellowtail flounder in the Canadian Maritimes region of the northwest Atlantic Ocean. Can. Tech. Rep. Fish. Aquat. Sci. 1196, 74 p.
- 1983e. Variability in size and age at sexual maturity of witch flounder, *Glyptocephalus cynoglossus*, in the Canadian Maritimes region of the northwest Atlantic Ocean. Can. Field-Nat. 97(4):409-422.
- Beacham, T.D., and S.J. Nepszy. 1980. Some aspects of the biology of white hake, *Urophycis tenuis*, in the southern Gulf of St. Lawrence. J. Northw. Atl. Fish. Sci. 1:49-54.
- Beverton, R.J.H. 1963. Maturation, growth and mortality of clupeid and engraulid stocks in relation to fishing. In Contrib. to hering symp. 1961 (B.B. Parrish, ed), p. 44-67. Rapp. P.-v. Reun. Cons. perm. int. Explor. Mer, 154.
1987. Longevity in fish: some ecological and evolutionary considerations. In Evolution of longevity in animals (A.D. Woodhead and K.H. Thompson, eds.), p. 161-186. Basic Life Sciences, Vol. 42. Plenum Press, NY.
- Beverton, R.J.H., and S.J. Holt. 1959. A review of the lifespans and mortality rates of fish in nature, and their relation to growth and other physiological characteristics. In CIBA Foundation colloquia on ageing. Vol. 5: the lifespan of animals (G.E.W. Wolstonholme, and O'Connor, eds.), p.142-180. Little, Brown and Co., Boston, 324 p.
- Bigelow, H.B. and W.C. Schroeder. 1953. Fishes of the Gulf of Maine. U.S. Fish Wildl. Serv. Fish. Bull. 74, 577 p.
- Bowering, W.R. 1976. Distribution, age and growth, and sexual maturity of witch flounder (*Glyptocephalus cynoglossus*) in Newfoundland waters. J. Fish. Res. Board Can. 33:1574-1584.
1987. Distribution of witch flounder, *Glyptocephalus cynoglossus*, in the southern Labrador and eastern Newfoundland area and changes in certain biological parameters after 20 years of exploitation. Fish. Bull., U.S. 85(3):611-629.
1989. Distribution of witch flounder, *Glyptocephalus cynoglossus*, in the southern Newfoundland area and changes in age, growth, and sexual maturity patterns since commercial exploitation began in the early 1950s. Trans. Am. Fish. Soc. 118:251-271.
- Bowering, W.R., and W.B. Brodie. 1984. Distribution of witch flounder in the northern Gulf of St. Lawrence and changes in its growth and sexual maturity patterns. N. Am. J. Fish. Manage. 4(4A):399-413.
- Burnett, J., M.R. Ross, and S.H. Clark. 1992. Several biological aspects of the witch flounder, *Glyptocephalus cynoglossus* (L.), in the Gulf of Maine-Georges Bank region. J. Northw. Atl. Fish. Sci. 12:15-25.
- Burnett, J., L. O'Brien, R.K. Mayo, J. Darde, and M. Bohan. 1989. Finfish maturity sampling and classification schemes used during Northeast Fisheries Center bottom trawl surveys, 1963-1989. NOAA Tech. Mem. NMFS-F/NEC-76, iii + 14 p.
- Burton, M.P., and D.R. Idler. 1984. The reproductive cycle in winter flounder, *Pseudopleuronectes americanus* (Walbaum). Can. J. Zool. 62:2563-2567.

- Cole, L.C.  
1954. The population consequences of life history phenomena. *Q. Rev. Biol.* 29:103-137.
- Dixon, W.J. (ed)  
1981. BMDP Statistical Software. Univ. Calif. Press, Berkeley, 725 p.
- Doubleday, W.G.  
1981. Manual on groundfish surveys in the northwest Atlantic. NAFO Scientific Council Studies No. 2, 55 p.
- Doubleday, W.G., and R.G. Halliday.  
1975. An analysis of the silver hake fishery on the Scotian Shelf. *Int. Comm. Northwest Atl. Fish., Res. Doc.* 75/104, Serial No. 3597, 34 p.
- DuPaul, W.D. and J.D. McEachran.  
1973. Age and growth of the butterfish, *Peprius triacanthus*, in the lower York River. *Chesapeake Sci.* 14(3):205-207.
- Elkin, H.W.  
1955. Whiting in Irish waters. *Proc. R. Dublin Soc. N.S.*, 27:17-30.
- Fahay, M.P., and K.W. Able.  
1989. White hake, *Urophycis tenuis*, in the Gulf of Maine: spawning seasonality, habitat use, and growth in young of the year and relationships to the Scotian Shelf population. *Can. J. Zool.* 67:1715-1724.
- Finkelstein, S.L.  
1969a. Age and growth of scup in the waters of Eastern Long Island. *New York Fish and Game J.* 16(1):84-110.  
1969b. Age at maturity of scup from New York waters. *New York Fish and Game J.* 16(2):224-237.
- Fogarty, M.J., G. Delaney, J.W. Gillikin, J.C. Poole, D.E. Ralph, P.G. Scarlett, R.W. Smith and S.J. Wilk.  
1983. Stock discrimination of summer flounder (*Paralichthys dentatus*) in the Middle and South Atlantic Bight: results of a workshop. NOAA Tech. Mem. NMFS-F/NEC-18, 14 p.
- Gabriel, W.L., M.P. Sissenwine, and W.J. Overholtz.  
1989. Analysis of spawning stock biomass per recruit: an example for Georges Bank haddock. *North Am. J. Fish. Manage.* 9(4):383-391.
- Garrod, D.J., and J.W. Horwood.  
1984. Reproductive strategies and the response to exploitation. In *Fish Reproduction: Strategies and tactics* (G.W. Potts and R.J. Wootton, eds.) p. 367-381. Acad. Press, London.
- Gillikin, J.W., Jr., and B.F. Holland.  
1983. Summary of summer flounder (*Paralichthys dentatus*) maturity data from the Atlantic Ocean offshore North Carolina. Rep., North Carolina Dept. of Natural Resources and Community Dev., Div. of Mar. Fish., 7 p.
- Godo, O.R., and E. Moksness.  
1987. Growth and maturation of Norwegian coastal cod and northeast Arctic cod under different conditions. *Fish. Res.* 5:235-242.
- Grosslein, M.D.  
1962. Haddock stocks in the ICNAF Convention area. *Int. Comm. Northwest Atl. Fish. Redb.* 1962, Part III:124-131.
- Halliday, R.G.  
1987. Size and age at sexual maturity of Atlantic argentine, *Argentina silus*: a critique. *Environ. Biol. Fishes* 19(2):139-147.
- Horn, M.H.  
1970. Systematics and biology of the stromateid fishes of the genus *Peprius*. *Bull. Mus. Comp. Zool.* 149(5):165-261.
- Howe, A.B. and P.G. Coates.  
1975. Winter flounder movements, growth, and mortality off Massachusetts. *Trans. Am. Fish. Soc.* 104:13-29.
- Johnson, L.L. and E. Casillas.  
1991. The use of plasma parameters to predict ovarian maturation state in English sole *Parophrys vetulus* Girard. *J. Exp. Mar. Biol. Ecol.* 151: 257-270.
- Jørgensen, T.  
1990. Long-term changes in age at sexual maturity of north-east Arctic cod (*Gadus morhua* L.). *J. Cons. int. Explor. Mer* 46:235-248.
- Keats, D.W., G.R. South, and D.H. Steele.  
1985. Reproduction and egg guarding by Atlantic wolffish (*Anarhichas lupus*: Anarhichidae) and ocean pout (*Macrozoarces americanus*: Zoarcidae) in Newfoundland waters. *Can. J. Zool.* 63:2565-2568.
- Kennedy, V.S., and D.H. Steele.  
1971. The winter flounder, (*Pseudopleuronectes americanus*) in Long Pond, Conception Bay, Newfoundland. *J. Fish. Res. Board Can.* 28:1153-1165.
- Lavenda, N.  
1949. Sexual differences and normal protogynous hermaphroditism in the Atlantic sea bass *Centropristis striata*. *Copeia* 1949(3):185-194.
- Livingstone, R., Jr., and L. Dery.  
1976. An observation on the age and length at maturity of cod in the Georges and Browns stocks. *Int. Comm. Northwest Atl. Fish. Res. Doc.* 76/VI/42. Serial No. 3826, 2 p.
- Lux, F.E.  
1963. Identification of New England yellowtail flounder groups. *Fish. Bull., U.S.* 63(1):1-10.  
1973. Age and growth of the winter flounder, *Pseudopleuronectes americanus*, on Georges Bank. *Fish. Bull., U.S.* 71(2):505-512.
- Lux, F.E., A.E. Peterson, and R.J. Hutton.  
1970. Geographical variation in fin ray number in winter flounder, *Pseudopleuronectes americanus* (Walbaum) off Massachusetts. *Trans. Am. Fish. Soc.* 99:483-488.
- MacKay, K. T.  
1967. An ecological study of mackerel *Scomber scombrus* (Linnaeus) in the coastal waters of Canada. *Fish. Res. Board Can., Tech. Rep. No.* 31. 127 p.  
1973. Aspects of the biology of Atlantic mackerel in ICNAF subarea 4. *Int. Comm. Northwest Atl. Fish. Res. Doc.* 73/70 Serial No. 3019, 11 p.
- Markle, D.F., D.A. Methven, and L.J. Coates-Markle.  
1982. Aspects of spatial and temporal co-occurrence in the life history stages of the sibling hakes, *Urophycis chuss* (Walbaum 1792) and *Urophycis tenuis* (Mitchill 1815) (Pisces: Gadidae). *Can. J. Zool.* 60:2057-2078.
- Mayo, R.K.  
1980. Exploitation and biological status of redfish, *Sebastes fasciatus* (L.), in the Gulf of Maine-Georges Bank region, with particular reference to the 1971 year-class. *J. Northw. Atl. Fish. Sci.* 1:21-37.
- Mayo, R.K., J. Burnett, T.D. Smith, and C.A. Muchant.  
1990. Growth-maturation interactions of Acadian redfish, (*Sebastes fasciatus* Storer) in the Gulf of Maine-Georges Bank region of the northwest Atlantic. *J. Cons. int. Explor. Mer* 46:287-305.
- Mayo, R.K., J.M. McGlade, and S.H. Clark.  
1989. Patterns of exploitation and biological status of pollock (*Pollachius virens* L.) in the Scotian Shelf, Georges Bank, and Gulf of Maine area. *J. Northw. Atl. Fish. Sci.* 9:13-36.
- Mercer, L.P.  
1978. The reproductive biology and population dynamics of black sea bass, *Centropristis striata*. Ph.D. diss., College of William and Mary, Williamsburg, VA, 196 p.

- Molander, A.R.  
1925. Observations on the witch flounder and its growth. Publ. Circ. Cons. Explor. Mer 85, 15 p.
- Moore, E.  
1947. Studies on the marine resource of Southern New England, VI. The sand flounder, *Lophopsetta aquosa*. Bull. Bingham Oceanogr. Coll. (Yale Univ.) XI, art. 3, 79 p.
- Moores, J.A., G.H. Winters, and L.S. Parsons.  
1975. Migrations and biological characteristics of Atlantic mackerel (*Scomber scombrus*) occurring in Newfoundland waters. J. Fish. Res. Bd. Can. 32:1347-1357.
- Morrison, C.M.  
1990. Histology of the Atlantic cod, *Gadus morhua*: an atlas, Part III. Reproductive tract. Can. Spec. Pub. Fish. Aquat. Sci. 110, 177 p.
- Morse, W.W.  
1981. Reproduction of the summer flounder, *Paralichthys dentatus* (L.). J. Fish. Biol. 19:189-203.
- Murawski, S.A., and G.T. Waring.  
1979. A population assessment of butterfish, *Peprilus triacanthus*, in the northwestern Atlantic Ocean. Trans. Am. Fish. Soc. 108:427-439.
- Musick, J.A.  
1969. The comparative biology of two American Atlantic hakes, *Urophycis chuss* and *U. tenuis* (Pisces, Gadidae). Ph.D. diss., Harvard Univ., Cambridge, MA, 150 p.
- New England Fishery Management Council.  
1985. Fishery Management Plan for the Northeast Multispecies Fishery. New England Fishery Management Council, Saugus, Massachusetts, 479 p.
- Ni, I-H., and E.J. Sandeman.  
1984. Size at maturity for northwest Atlantic redfishes (*Sebastes*). Can. J. Fish. Aquat. Sci. 41:1753-1762.
- NEFC (Northeast Fisheries Center).  
1988. An evaluation of the bottom trawl survey program of the Northeast Fisheries Center. NOAA Tech. Mem. NMFS-F/NEC-52, 83 p.
- NEFSC (Northeast Fisheries Science Center).  
1991. Status of the Fishery Resources off the northeastern United States for 1990. NOAA Tech. Mem., NMFS-F/NEC-81, 132 p.
- O'Brien, L.  
1990. Effects of fluctuations in stock abundance upon life history parameters of Atlantic cod, *Gadus morhua* L., for the 1970-1987 year classes from Georges Bank and the Gulf of Maine. M. Sci. thesis, University of Washington, Seattle, Wa., 95 p.
- Olsen, Y. H., and D. Merriman.  
1946. Studies on the marine resources of Southern New England. IV. The biology and economic importance of the ocean pout, *Macrozoarces americanus* (Bloch and Schneider). Bull. Bingham Oceanogr. Coll. (Yale Univ.) 9(4), 184 p.
- Overholtz, W. J.  
1987. Factors relating to the reproductive biology of Georges Bank haddock (*Melanogrammus aeglefinus*) in 1977-83. J. Northw. Atl. Fish. Sci. 7:145-154.  
1989. Density-dependent growth in the northwest Atlantic stock of Atlantic mackerel (*Scomber scombrus*). J. Northw. Atl. Fish. Sci. 9:115-121.
- Penttilä, J. and L. M. Dery (eds).  
1988. Age determination methods for northwest Atlantic species. NOAA Tech. Rep. NMFS 72, 135 p.
- Penttilä, J., G.A. Nelson, and J.M. Burnett.  
1989. Guidelines for estimating lengths at age for 18 northwest Atlantic finfish and shellfish species. NOAA. Tech. Mem. NMFS-F/NEC-66, iii + 39 p.
- Perlmutter, A.  
1947. The blackback flounder and its fishery in New England and New York. Bull. Bingham Oceanogr. Coll., (Yale Univ.) 11(2):1-92.
- Perlmutter, A., and G.M. Clarke.  
1949. Age and growth of immature rosefish (*Sebastes marinus*) in the Gulf of Maine and off Nova Scotia. Fish. Bull., U.S. 51:207-228.
- Pierce, D.E., and A.B. Howe.  
1977. A further study on winter flounder group identification off Massachusetts. Trans. Am. Fish. Soc. 106:131-139.
- Roff, D.A.  
1981. On being the right size. Am. Nat. 118:405-422.  
1982. Reproductive strategies in flatfish: a first synthesis. Can. J. Fish. Aquat. Sci. 39:1686-1698.
- Royce, W.F., R. J. Buller, and E.D. Premetz.  
1959. Decline of the yellowtail flounder (*Limanda ferruginea*) off New England. Fish. Bull., U.S. 146, 59:169-267.
- Schaffer, W.M., and P.F. Elson.  
1975. The adaptive significance of variations in life history among local populations of Atlantic salmon in North America. Ecology 56:577-590.
- Scott, D. M.  
1954. A comparative study of the yellowtail flounder from three Atlantic fishing areas. J. Fish. Res. Bd. Can. 11(3):171-197.
- Sette, O.E.  
1950. Biology of the Atlantic mackerel (*Scomber scombrus*). Part II. Migrations and habits. U.S. Fish and Wildlife Ser. Fish. Bull. 49(51):251-358.
- Shepherd, G.R.  
1991. Meristic and morphometric variation in black sea bass north of Cape Hatteras, North Carolina. North Am. J. Fish. Manage. 11(2):139-148.
- Sinclair, A., M. Sinclair, and T.D. Iles.  
1980. An analysis of some biological characteristics of the 4X juvenile herring fishery. Can. Atl. Fish. Sci. Adv. Committee Res. Doc. 80/20, 28 p.
- Sinclair, M., J. Black, T.D. Iles, and W. Stobo.  
1979. Preliminary analysis of the use of Bay of Fundy larval survey data in 4WX herring assessments. Can. Atl. Fish. Sci. Adv. Committee Res. Doc. 79/44, 18 p.
- Sinclair, M., A. Sinclair, and T.D. Iles.  
1982. Growth and maturation of southwest Nova Scotia Atlantic herring (*Clupea harengus harengus*). Can. J. Fish. Aquat. Sci. 39:288-295.
- Sissenwine, M.P. and J.G. Shepherd.  
1987. An alternative perspective on recruitment overfishing and biological reference points. Can. J. Fish. Aquat. Sci. 44:913-918.
- Smith, R.W., L.M. Dery, P.G. Scarlett, and A. Jearld Jr.  
1981. Proceedings of the summer flounder (*Paralichthys dentatus*) age and growth workshop; 20-21 May 1980, Northeast Fisheries Center, Woods Hole, MA. NOAA Tech. Mem. NMFS-F/NEC-11, 30 p.
- Steele, D.H.  
1963. Pollock (*Pollachius virens* (L.)) in the Bay of Fundy. J. Fish. Res. Bd. Can. 20(5): 1267-1314.
- Sullivan, L.F.  
1981. American plaice, *Hippoglossoides platessoides* in the Gulf of Maine: I. The fishery II. Age and growth III. Spawning and larval distribution. M. Sci. thesis. University of Rhode Island, Kingston, R.I., 95 p.
- Szedlmayer, S.T., K.W. Able, and R.A. Roundtree.  
1992. Growth and temperature - induced mortality of young-of-the-year summer flounder, *Paralichthys dentatus*, in southern New Jersey. Copeia 1992(1):120-128.

- Templeman, W., and C.A. Bishop.  
1979. Sexual maturity and spawning in haddock, *Melanogrammus aeglefinus*, of St. Pierre Bank. Int. Comm. Northwest Atl. Fish., Res. Bull. 14:77-83.
- Templeman, W., V.M. Hodder, and R. Wells.  
1978. Sexual maturity and spawning in haddock, *Melanogrammus aeglefinus*, of the southern Grand Bank. Int. Comm. Northwest Atl. Fish. Res. Bull. 13:53-63.
- Thorpe, L.A.  
1991. Aspects of the biology of the windowpane flounder, *Scophthalmus aquosus*, in the northwest Atlantic Ocean. M. Sci. thesis, Dept. of Forestry and Wildlife, Univ. of Massachusetts-Amherst, Amherst, MA., 85 p.
- Wenner, C.A., W.A. Roumillat, and C.W. Waltz.  
1986. Contributions to the life history of black sea bass, *Centropristis striata*, off the southeastern United States. Fish. Bull., U.S. 84(3):723-741.
- Winters, G.H.  
1976. Recruitment mechanisms of southern Gulf of St. Lawrence Atlantic herring (*Clupea harengus harengus*). J. Fish. Res. Board Can. 33:1751-1763.
- Wise, J.P.  
1963. Cod groups in the New England area. Fish Bull., U.S. 63(1):189-203.
- Zar, J.H.  
1984. Biostatistical analysis. Prentice-Hall, Inc., Englewood Cliffs, 718 p.





## *Guide for Authors*

### NOAA Technical Report NMFS

#### **PREPARATION**

---

**Title** of the paper should be as brief as possible (between 6 and 12 words) and should not include the Latin binomial except when it is needed for clarification, i.e., to distinguish between species sharing the same common name or between species known internationally by different common names.

**Abstract** should be one paragraph of about 200 words or less. It should state the main scope of the paper but emphasize its conclusions and relevant findings. Because abstracts are circulated by abstracting agencies, it is important that they represent the research clearly and concisely.

**Text** must be typed double-spaced throughout and should be divided into the following sections: **Introduction**, **Materials and Methods**, **Results**, **Discussion** (or **Conclusions**), and **Acknowledgments**. Headings within each section must be short, reflect a logical sequence, and follow the rules of multiple subdivision (i.e., there can be no subdivision without at least two items of subdivision). The entire text should be intelligible to interdisciplinary readers; therefore, all acronyms, abbreviations, and technical terms should be defined the first time they are used. The scientific names of species must be written in full the first time they are mentioned and abbreviated in subsequent references. Footnotes should be avoided as much as possible. We follow the *U.S. Government Printing Office, Style Manual* (1984 ed.) and the *CBE Style Manual* (5th ed.) for general format and style; the American Fisheries Society's most recent edition of *Common and Scientific Names of Fishes from the United States and Canada* for fish nomenclature. Dates should be written as 11 November 1991. Measurements should be expressed in metric units, e.g., tons as metric tons (t), but if the work is in British long tonnes, please make this fact explicit to the reader. The numeral one (1) should be typed as a one, not a lower-case el (l).

**Citations** comprise both unpublished and published works. Authors are advised to avoid references to nonstandard material such as internal and project reports wherever possible. For these works, include whether they are available from NTIS (National Technical Information Service) or from some other public depository. Personal communications and unpublished data must be cited in parentheses in the text with full address of communicator or author. Follow the name-and-year system for citation format. In the text, cite Smith and Jones (1977) or (Smith and Jones 1977). If there is a sequence of citations, list chronologically: Smith 1932; Green 1947; Smith and Jones 1985. Abbreviations of serials should conform to abbreviations given in *Serial Sources for Biosis Data Base*<sup>TM</sup>. **AUTHORS ARE RESPONSIBLE FOR THE ACCURACY AND COMPLETENESS OF ALL CITATIONS.**

**Tables** should not be excessive in size and must be cited in numerical order in the text. All unusual symbols must be explained in the table heading. Other incidental comments can be

footnoted with italic lower case letters (*a, b, c*) for general comment, with asterisks for probability in statistical data. Because tables are typeset, they need only be submitted typed and formatted.

**Figures** include line illustrations and photographs. Unless photographs are submitted on glossy paper with good contrast, we cannot guarantee a good final printed copy. Figures must be labeled with author's name and number of figure. Lettering within the figure should not be so heavy and large as to upstage the impact of the entire figure. Avoid "outlier" words in a figure which take up needless space. Each figure must include a legend that explains all symbols and abbreviations. These figure legends should be typewritten on a separate piece of paper at the end of the manuscript.

**Copies of Published Reports** are available free of charge to the senior author (50 copies) and to his or her laboratory (50). If the article is part of a proceedings or collection of articles, the senior author receives 50 free copies of the individual article and one copy of the collective work. Additional copies (reprints or offprints) of an article may be purchased in lots of 100 at a cost of \$1.40 per page at the time of first printing. Additional copies of a collective work must be sought from the editor assigned to coordinate the research papers for submission as a NOAA Technical Report.

#### **SUBMISSION**

---

Send manuscript (original and two copies) to the Scientific Editor:

Dr. Ronald Hardy, Scientific Editor  
Northwest Fisheries Science Center, F/NWC3  
National Marine Fisheries Service, NOAA  
2725 Montlake Boulevard East  
Seattle, WA 98112.

Once the manuscript has been accepted for publication, you will be asked to submit a software copy of your manuscript to the Scientific Editor. The software copy should be submitted in ASCII format (i.e., in a MS-DOS "print" or "nondocument" file) and should be placed on a 5.25-inch (preferably) or 3.5-inch disk that is double-sided, double or high density, and that is compatible with either IBM or Apple Macintosh systems.

Once the manuscript is being typeset and prepared for publication, all inquiries should be made to the Managing Editor:

Sharyn Matriotti, Managing Editor  
NOAA Technical Reports NMFS  
National Marine Fisheries Service  
Scientific Publications Office, F/NWR1  
7600 Sand Point Way NE  
Seattle, WA 98115-0070.

UNITED STATES  
DEPARTMENT OF COMMERCE

NATIONAL OCEANIC AND ATMOSPHERIC  
ADMINISTRATION  
NATIONAL MARINE FISHERIES SERVICE  
SCIENTIFIC PUBLICATIONS OFFICE  
BIN C15700  
SEATTLE, WA 98115  
OFFICIAL BUSINESS

Penalty for Private Use, \$300

BULK RATE  
POSTAGE & FEES PAID  
U.S. Department of Commerce  
Permit No. G-19

## NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

*The National Oceanic and Atmospheric Administration* was established as part of the Department of Commerce on October 13, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

**PROFESSIONAL PAPERS**—Important definitive research results, major techniques, and special investigations.

**CONTRACT AND GRANT REPORTS**—Reports prepared by contractors or grantees under NOAA sponsorship.

**ATLAS**—Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

**TECHNICAL SERVICE PUBLICATIONS**—Reports containing data, observations, instructions, etc. A partial listing includes data serials; predictions and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

**TECHNICAL REPORTS**—Journal quality with extensive details, mathematical developments, or data listings.

**TECHNICAL MEMORANDUMS**—Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



*Information on availability of NOAA publications can be obtained from:*

U.S. Department of Commerce  
National Technical Information Service  
5285 Port Royal Road  
Springfield, VA 22161